
# CHAMOT

Recombinant IL-1 beta, Human, GMP

CM002-100HPG

CM002-1000HPG





- 1 Product Specifications
- 2 Product Note
- 3 Storage/Shipping
- 4 Scientific Data



## **Product Datasheet**

# Recombinant IL-1 beta, Human, GMP

Catalog#	CM002-100HPG	CM002-1000HPG
Size	100 ug	1 mg

### **Product Specifications**

Background	Interleukin-1 beta (IL-1 $\beta$ ) is a major cytokine expressed in macrophage, NK cells, monocytes, and neutrophils. It also plays fundamental role in inflammatory response, including monocyte activation, which is essential for the host defense and pathogen and pathogen resistance. Pro-IL-1 $\beta$ is cleaved by cytosolic caspase 1 to form mature IL-1 $\beta$ .
Synonyms	interleukin 1 beta, Catabolin, Lymphocyte-Activating Factor (LAF), Endogenous Pyrogen (EP), Leukocyte Endoge_x005f_x005f_x005f_nous Mediator (LEM), Mononuclear Cell Factor (MCF), IL1, IL1B
Uniprot ID	P01584
Molecular Weight	The protein has a calculated MW of 18.5 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).
Expression System	Escherichia coli
Purity	>98% as determined by SDS-PAGE analysis.
	> 30% as determined by 3D3-FAGE analysis.
Activity	Measure by its ability to induce IL-8 secretion in HT29 cells. The ED <sub>50</sub> for this effect is <2.0 ng/mL. The specific activity of recombinant human IL-1 beta is approximately >3 x 10 <sup>7</sup> IU/mg, which is calibrated against the human IL-1 beta WHO International Standard (NIBSC code: 86/680).
·	Measure by its ability to induce IL-8 secretion in HT29 cells. The ED $_{50}$ for this effect is <2.0 ng/mL. The specific activity of recombinant human IL-1 beta is approximately >3 x $10^7$ IU/mg, which is calibrated against the human IL-1 beta WHO International Standard (NIBSC
Activity Endotoxin	Measure by its ability to induce IL-8 secretion in HT29 cells. The ED $_{50}$ for this effect is <2.0 ng/mL. The specific activity of recombinant human IL-1 beta is approximately >3 x 10 $^7$ IU/mg, which is calibrated against the human IL-1 beta WHO International Standard (NIBSC code: 86/680).
Activity  Endotoxin Level  Protein	Measure by its ability to induce IL-8 secretion in HT29 cells. The ED <sub>50</sub> for this effect is <2.0 ng/mL. The specific activity of recombinant human IL-1 beta is approximately >3 x 10 <sup>7</sup> IU/mg, which is calibrated against the human IL-1 beta WHO International Standard (NIBSC code: 86/680).  <0.05 EU per 1 µg of the protein by the LAL method. MASAPVRSLNCTLRDSQQKSLVMSGPYELKALHLQGQDMEQQVVFSMSFVQGEESNDKIPVALGLKE KNLYLSCVLKDDKPTLQLESVDPKNYPKKKMEKRFVFNKIEINNKLEFESAQFPNWYISTSQAENMPVFL

#### **Product Note**

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration not less than 0.5 mg/mL and incubate the stock solution at RT for at least 20 min to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

#### Storage/Shipping


Stability & Storage


Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.


**Shipping** 

Blue Ice

#### **Scientific Data**







GMP Human IL-1 beta Protein, His Tag, E. coli induces IL-8 secretion in HT29 cells, with the ED<sub>50</sub> < 2.0

ng/mL.

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491