

CHAMOT

**重组人白细胞介素7 / Recombinant IL-7, Human,
GMP**

CM009-100HPG

CM009-1000HPG



CHAMOT

乔默®生物

Specialize In Cytokines



CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组人白细胞介素7 / Recombinant IL-7, Human, GMP

产品编号	CM009-100HPG	CM009-1000HPG
规格	100 ug	1 mg

产品简介

背景描述 Interleukin-7 (IL-7) is a multipotent cytokine belonged to one of the members of IL-2 superfamily. IL-7 has diverse effects on the hematopoietic and immune regulations. IL-7 is a trophic factor that is necessary for both B cell and T cell proliferation and development. In addition, it presents potential antitumor effects in tumors such as glioma, melanoma, lymphoma, leukemia, prostate cancer, and glioblastoma.

别称 interleukin 7, Lymphopoietin 1(LP-1), pre-B-cell factor, IL7

蛋白编码 P13232

分子量 The protein has a calculated MW of 18.3 kDa. The protein migrates as 19 kDa under reducing condition (SDS-PAGE analysis).

表达系统 Escherichia coli

纯度 >97% as determined by SDS-PAGE analysis.

生物活性 Measured by its ability to induce PHA-activated human PBMCs proliferation. The ED₅₀ for this effect is <0.8 ng/mL. The specific activity of recombinant human IL-7 GMP is > 1 x 10⁸ unit/mg, which is calibrated against the human IL-7 WHO Reference Reagent (NIBSC code: 90/530).

内毒素检测 <0.05 EU per 1 µg of the protein by the LAL method.

蛋白序列 MDCDIEGKDGKQYESVLMVSIQLLDSMKEIGSNCLNNEFNFFKRHICDANKEGMFLFRAAR KLRQFLKMNSTGDFDLHLLKVSEGTILLNCTGQVKGRKPAALGEAQPTKSLEENKSLKEQKKL NDLCFLKRLQEIKTCWNKILMGTKEH with polyhistidine tag at the C-terminus.

支原体 Not detected

产品形式 Lyophilized from a 0.2 µm filtered solution of PBS, pH 8.0.

产品应用 Cell Culture


产品使用

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 0.5 mg/mL and incubate the stock solution at RT for at least 20 min to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

产品储存/运输

储存	Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.
运输	Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491