# CHAMOT

# Recombinant IFN gamma, Human, GMP

CM081-100HPG

CM081-1000HPG





- 1 Product Specifications
- 2 Product Note
- 3 Storage/Shipping
- 4 Scientific Data



## **Product Datasheet**

# Recombinant IFN gamma, Human, GMP

Catalog#	CM081-100HPG	CM081-1000HPG
Size	100 ug	1 mg

### **Product Specifications**

Background	The cytokine IFN gamma could protect cells from viral infections and belongs to the family of interferons. A lot of studies have shown that IFN gamma secreted by antigen triggered cell types, including T cells, naive CD4+ T cells, macrophages, dendritic cells, and B cells. IFN gamma plays an important role to trigger the macrophage act against a diverse group of microbial targets, and the pleiotropic molecule associated with antiproliferative, proapoptotic and antitumor mechanisms. Based on the effector cytokine considered as a major effector of immunity, it has been used in the treatment of several diseases.
Synonyms	interferon gamma, Type II interferon, T-cell interferon, MAF
Uniprot ID	P01579
Molecular Weight	The protein has a calculated MW of 17.7 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).
Expression System	Escherichia coli
Purity	>95% as determined by SDS-PAGE analysis.
Activity	Measure by its ability to induce cytotoxicity in HT29 cells. The ED $_{50}$ for this effect is <1 ng/mL. The specific activity of recombinant human IFN gamma is approximately >2 x $10^6$ IU/mg, which is calibrated against the human IFN Gamma WHO Reference Material (NIBSC code: 87/586).
Activity Endotoxin Level	ng/mL. The specific activity of recombinant human IFN gamma is approximately >2 x 10 <sup>6</sup> IU/mg, which is calibrated against the human IFN Gamma WHO Reference Material (NIBSC
Endotoxin	ng/mL. The specific activity of recombinant human IFN gamma is approximately $>2 \times 10^6$ IU/mg, which is calibrated against the human IFN Gamma WHO Reference Material (NIBSC code: 87/586).
Endotoxin Level Protein	ng/mL. The specific activity of recombinant human IFN gamma is approximately >2 x 10 <sup>6</sup> IU/mg, which is calibrated against the human IFN Gamma WHO Reference Material (NIBSC code: 87/586).  <0.05 EU per 1 µg of the protein by the LAL method.  MQDPYVKEAENLKKYFNAGHSDVADNGTLFLGILKNWKEESDRKIMQSQIVSFYFKLFKNFKDDQSIQ KSVETIKEDMNVKFFNSNKKKRDDFEKLTNYSVTDLNVQRKAIHELIQVMAELSPAAKTGKRKRSQMLF

#### **Product Note**

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration not less than 0.5 mg/mL and incubate the stock solution at RT for at least 20 min to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

#### Storage/Shipping


Stability & Storage


Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

**Shipping** 

Blue Ice

#### **Scientific Data**





SDS- PAGE analysis of GMP human IFN gamma

GMP Human IFN Gamma Protein, His Tag, E. coli induces cytotoxicity in HT29 cells, with the ED $_{50}$  <1.0

ng/mL.

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491