

CHAMOT

Recombinant IFN gamma, Human, GMP

CM081-100HPG

CM081-1000HPG



CHAMOT

乔默®生物

Specialize In Cytokines



CONTENT

1 Product Specifications

2 Product Note

3 Storage/Shipping

4 Scientific Data

Recombinant IFN gamma, Human, GMP

Catalog#	CM081-100HPG	CM081-1000HPG
Size	100 ug	1 mg

Product Specifications

Background	The cytokine IFN gamma could protect cells from viral infections and belongs to the family of interferons. A lot of studies have shown that IFN gamma secreted by antigen triggered cell types, including T cells, naive CD4+ T cells, macrophages, dendritic cells, and B cells. IFN gamma plays an important role to trigger the macrophage act against a diverse group of microbial targets, and the pleiotropic molecule associated with antiproliferative, pro-apoptotic and antitumor mechanisms. Based on the effector cytokine considered as a major effector of immunity, it has been used in the treatment of several diseases.
Synonyms	interferon gamma, Type II interferon, T-cell interferon, MAF
Uniprot ID	P01579
Molecular Weight	The protein has a calculated MW of 17.7 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).
Expression System	Escherichia coli
Purity	>95% as determined by SDS-PAGE analysis.
Activity	Measure by its ability to induce cytotoxicity in HT29 cells. The ED ₅₀ for this effect is <1 ng/mL. The specific activity of recombinant human IFN gamma is approximately >2 x 10 ⁶ IU/mg, which is calibrated against the human IFN Gamma WHO Reference Material (NIBSC code: 87/586).
Endotoxin Level	<0.05 EU per 1 µg of the protein by the LAL method.
Protein Sequence	MQDPYVKEAENLKKYFNAGHSDVADNGTLFLGILKNWKEESDRKIMQSQIVSFYFKLFKNFKDDQSIQ KSVETIKEDMNVKFFNSNKKKRDDFEKLNTNYSVTDLNVQRKAIHELIQVMAELSPAAGTGKRKRSQMLF QGRRASQ with polyhistidine tag at the C-terminus.
Form	Lyophilized from a 0.2 µm filtered solution of PBS, pH 8.0.
Application	Cell Culture

Product Note

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 0.5 mg/mL and incubate the stock solution at RT for at least 20 min to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Storage/Shipping


Stability & Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.


Shipping

Blue Ice

Scientific Data



SDS- PAGE analysis of GMP human IFN gamma



GMP Human IFN Gamma Protein, His Tag, E. coli induces cytotoxicity in HT29 cells, with the ED₅₀ <1.0 ng/mL.