
CHAMOT

Recombinant Activin A, Human, GMP

CM118-100HPG

CM118-1000HPG





トスツと

- 1 Product Specifications
- 2 Product Note
- 3 Storage/Shipping
- 4 Scientific Data



Product Datasheet

Recombinant Activin A, Human, GMP

Catalog#	CM118-100HPG	CM118-1000HPG
Size	100 ug	1 mg

Product Specifications

Background	Activin is the member of the TGF beta superfamily of cytokines. The current understanding of Activin A is classified as a hormone, a growth factor, and a cytokine. Depending on the different cell type, that overexpression of Activin A can either inhibit or promote cells division. There are important implications for tumor biology. Activin A also increase joint inflammation in rheumatoid arthritis (RA) and induces pathogenic mechanisms in the respiratory system.
Synonyms	Inhibin beta A chain, Activin beta-A chain, Erythroid differentiation protein
Uniprot ID	P08476
Molecular Weight	The protein has a calculated MW of 12.96 kDa. The protein migrates as 13-17 kDa under reducing condition (SDS-PAGE analysis).
Expression System	Escherichia coli
Purity	>97% as determined by SDS-PAGE analysis.
Activity	Measure by its ability to inhibit the proliferation of mouse MPC-11 cells. The ED $_{50}$ for this effect is <10 ng/mL. The specific activity of recombinant human Activin A is approximately >1.0 x 10^3 IU/mg.
Endotoxin Level	<0.05 EU per 1 µg of the protein by the LAL method.
Protein Sequence	GLECDGKVNICCKKQFFVSFKDIGWNDWIIAPSGYHANYCEGECPSHIAGTSGSSLSFHSTV INHYRMRGHSPFANLKSCCVPTKLRPMSMLYYDDGQNIIKKDIQNMIVEECGCS
Form	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Application	Cell Culture

Product Note

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 0.5 mg/mL and incubate the stock solution at RT for at least 20 min to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Storage/Shipping


Stability & Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

Shipping

Blue Ice

Scientific Data



SDS- PAGE analysis of GMP human Activin A

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491