
# **CHAMOT**

# Recombinant TPO, Human, GMP

CM124-100HPG

CM124-1000HPG





- 1 Product Specifications
- 2 Product Note
- 3 Storage/Shipping
- 4 Scientific Data



# **Product Datasheet**

# Recombinant TPO, Human, GMP

Catalog#	CM124-100HPG	CM124-1000HPG
Size	100 ug	1 mg

## **Product Specifications**

Background	Thrombopoietin (TPO) is a glycoprotein that produced by the liver, kidney, marrow stroma and several other tissues. The TPO level in the blood is mostly negatively correlated with the abundance of platelets and bone marrow megakaryocytes, although multiple states of inflammation or infection, liver failure, and hematological disturbances are associated with unexpectedly high or low circulating levels of the hormone.
Synonyms	thrombopoietin, Megakaryocyte colony-stimulating factor, c-MPL Ligand, MGDF, THPO
Uniprot ID	P40225
Molecular Weight	The protein has a calculated MW of 19.6 kDa. The protein migrates as 20 kDa under reducing condition (SDS-PAGE analysis).
Expression System	Escherichia coli
Purity	>95% as determined by SDS-PAGE analysis.
Activity	Measure by its ability to induce proliferation in MO7e cells. The ED $_{50}$ for this effect is <2 ng/mL.
Endotoxin Level	<0.05 EU per 1 µg of the protein by the LAL method.
Protein Sequence	SPAPPACDLRVLSKLLRDSHVLHSRLSQCPEVHPLPTPVLLPAVDFSLGEWKTQMEETKAQDILGAVTLLL EGVMAARGQLGPTCLSSLLGQLSGQVRLLLGALQSLLGTQLPPQGRTTAHKDPNAIFLSFQHLLRGKVR FLMLVGGSTLCVRRAPPTTAVPSRTSLVLTLNEL with polyhistidine tag at the N-terminus.
Form	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Application	Cell Culture

### **Product Note**

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration not less than 0.5 mg/mL and incubate the stock solution at RT for at least 20 min

to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

### Storage/Shipping


Stability & Storage


Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

**Shipping** 

Blue Ice

### **Scientific Data**





SDS- PAGE analysis of GMP human TPO

GMP Human TPO Protein, His Tag, E. coli induced MO7e cell proliferation.

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491