
# **CHAMOT**

# Recombinant SCF, Human, GMP

CM171-100HPG

CM171-1000HPG





# トスツと

- 1 Product Specifications
- 2 Product Note
- 3 Storage/Shipping
- 4 Scientific Data



# **Product Datasheet**

# Recombinant SCF, Human, GMP

Catalog#	CM171-100HPG	CM171-1000HPG
Size	100 ug	1 mg

## **Product Specifications**

Background
------------

Stem Cell Factor (SCF) is a 166-amino-acid protein with a monomeric molecular weight of approximately 18.5 kDa. SCF is the ligand for the tyrosine kinase receptor c-kit, which regulates other growth factors, such as granulocyte colony-stimulating factor (G-CSF), granulocyte macrophage-colony-stimulating factor (GM-CSF), and interleukin-3 to stimulate proliferation, and differentiation of hematopoietic stem cells. SCF can be produced by a variety of cells, including fibroblasts, smooth muscle, endothelial cells, mast cells and bone marrow macrophages endothelial cells.

	macrophages endothelial cells.
Synonyms	DCUA, DFNA69, FPH2, FPHH, KL-1, Kitl, MGF, SF, SHEP7, SLF, KIT ligand
Uniprot ID	P21583
Molecular Weight	The protein has a calculated MW of 19.4 kDa. The protein migrates as 21 kDa under reducing condition (SDS-PAGE analysis).
Expression System	Escherichia coli
Purity	>98% as determined by SDS-PAGE analysis.
Activity	Measure by its ability to induce TF-1 cells proliferation. The ED <sub>50</sub> for this effect is

Expression System	Escherichia coli
Purity	>98% as determined by SDS-PAGE analysis.
Activity	Measure by its ability to induce TF-1 cells proliferation. The ED $_{50}$ for this effect is <5 ng/mL. The specific activity of recombinant human SCF is > 5 x 10 $^{5}$ IU/mg.
Endotoxin Level	<0.05 EU per 1 $\mu$ g of the protein by the LAL method.
Protein Sequence	MEGICRNRVTNNVKDVTKLVANLPKDYMITLKYVPGMDVLPSHCWISEMVVQLSDSLTDL LDKFSNISEGLSNYSIIDKLVNIVDDLVECVKENSSKDLKKSFKSPEPRLFTPEEFFRIFNRSIDA FKDFVVASETSDCVVSSTLSPEKDSRVSVTKPFMLPPVA with polyhistidine tag at the Cterminus
Form	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Application	Cell Culture

### **Product Note**

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile  $H_2O$  to a concentration not less than 0.5 mg/mL and incubate the stock solution at RT for at least 20 min to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

### Storage/Shipping


Stability & Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

Shipping

Blue Ice

### **Scientific Data**



SDS- PAGE analysis of GMP human SCF

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491