


CHAMOT

重组人4-1BB配体 /Recombinant 4-1BBL, Human, Animal-Free

CM085-5HP



CM085-20HP




CM085-100HP



CM085-500HP



CM085-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENTS

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组人4-1BB配体 /Recombinant 4-1BBL, Human, Animal-Free

产品编号	CM085-5HP	CM085-20HP	CM085-100HP	CM085-500HP	CM085-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

4-1BB ligand (4-1BBL) is a type II transmembrane protein that is part of the tumor necrosis factor (TNF) ligand family. As an inducible co-stimulatory molecule, it presents on several antigen presenting cell (APC) types, including B cells, macrophages and DCs. The interactions between 4-1BB and 4-1BBL trigger pleiotropic effects on the immune response including antigen presenting process, proliferation of CD4 and CD8 positive T-cells, as well as cytokine secretion from T-cells through NF κ B, c-Jun, and p38 downstream signal pathways activation. Therefore, 4-1BB and 4-1BBL are recently used for the immunotherapy of cancer.

背景描述

别称

4-1BB ligand, CD137L, TNLG5A, TNFSF9, Tumor necrosis factor ligand superfamily member 9

蛋白编码

P41273

分子量

The protein has a calculated MW of 20.4 kDa. The protein migrates as 20 kDa under reducing condition (SDS-PAGE analysis).

表达系统

Escherichia coli

纯度

>95% as determined by SDS-PAGE.

生物活性

Measure by its ability to induce IL-8 secretion in human PBMCs. The ED₅₀ for this effect is 1-5 ng/mL.

内毒素检测

<0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列

MREGPELSPDDPAGLLDLRQGMFAQVQNVLLIDGPLSWYSDPGLAGVSLTGGLSYKEDTKE
 LVVAKAGVYYVFFQLELRRVVAGEGSGSVSLALHLQLPLRSAAGAAALALTVDLPPASSEARNSA
 FGFQGRLLHLSAGQRLGVHLHTEARARHAWQLTQGATVLGLFRVTPEIPAGLPSPRSE with
 polyhistidine tag at the C-terminus.

产品形式

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.
Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ($\geq 20 \mu\text{l}$).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant human 4-1BBL

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491