


CHAMOT

重组人B细胞激活因子 /Recombinant BAFF,
Human, Animal-Free

CM050-5HP



CM050-20HP





CM050-100HP



CM050-500HP



CM050-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1

产品简介

2

产品使用

3

产品储存/运输

4

实验数据展示

重组人B细胞激活因子 /Recombinant BAFF, Human, Animal-Free

产品编号	CM050-5HP	CM050-20HP	CM050-100HP	CM050-500HP	CM050-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

BAFF also known as BLYS, TALL-1 and TNFSF13B, which belongs to tumor necrosis factor family. BAFF is a 31.2 kDa type II transmembrane protein containing 285 residues that predominantly produced by myeloid cells. BAFF has been demonstrated to activate the survival of B-cells and the B-cell response by binding to BAFFR/BR3. Additionally, BAFF also takes part in regulating B and T cell function via forming two ligands-two receptors pathway through sharing TNFRSF13B/TACI and TNFRSF17/BCMA receptors with APRIL.

别称	B-cell activating factor, TNFSF13B, BLYS, CD257, DTL, TALL-1, TALL1, THANK, TNFSF20, TNLG7A, ZTNF4
蛋白编码	Q9Y275
分子量	The protein has a calculated MW of 17.98 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce IL-8 secretion in human PBMCs. The ED ₅₀ for this effect is <0.5 ng/mL.
内毒素检测	<0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列	MAVQGPEETVTQDCLQLIADSETPTIQKGSYTFVPWLLSFKRGSALEEKENKILVKETGYFFIYGQVLYTDKTYAMGHЛИQRKKVHVFGDELSVTLFRCIQNMPETLPNNSCYSAGIAKLEEGDELQLAIPRENAQISLDGDVTFFGALKLL with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ($\geq 20 \mu\text{l}$).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant human BAFF

Human BAFF, His Tag, E. coli induced IL-8 secretion in human PBMCs, with the ED₅₀ at 0.4031 ng/mL.

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491