


CHAMOT

重组人神经生长因子 β /Recombinant beta-NGF,
Human, Animal-Free

CM141-20HP




CM141-100HP



CM141-500HP



CM141-1000HP



CHAMOT

乔默[®]生物

Specialize In Cytokines

CONTENTS

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组人神经生长因子β /Recombinant beta-NGF, Human, Animal-Free

产品编号	CM141-20HP	CM141-100HP	CM141-500HP	CM141-1000HP
规格	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述 Nerve Growth Factors (NGF) is critical for the development and maintenance of the sympathetic and sensory neuron systems. NGF has been demonstrated as a complex that consists of three polypeptides named α , β and γ subunits. Among them, β subunit, which is known as beta-NGF, is a 26.9 kDa protein containing 241 residues that are involved in neuronal survival and differentiation. Besides, beta-NGF also acts as a ligand to TRKA receptor, which is indispensable for the differentiation and development of pain and temperature sensing neurons.

别称	nerve growth factor-beta, β -Nerve Growth Factor, NGF- β
蛋白编码	P01138
分子量	The protein has a calculated MW of 14.43 kDa. The protein migrates as 11 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Measured by its ability to induce TF-1 cells proliferation. The ED ₅₀ for this effect is <0.7 ng/mL. The specific activity of recombinant human beta-NGF is > 1 x 10 ⁶ IU/mg.
内毒素检测	<0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列	MSSSHPIFHRGEFSVCDSVSVWVGDKTTATDIKGKEVMVLGEVNINNSVFKQYFFETKCRDPN PVDSGCRGIDSKHWN SYCTTHTFKALTMDGKQAAWRFIRIDTACVCVLSRKAVRRA with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate, 0.2 M NaCl, pH 4.5. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ($\geq 20 \mu\text{l}$).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant human beta-
NGF

Human beta-NGF, His Tag, E. coli induced TF-1 cell proliferation,
with the ED_{50} at 0.6093 ng/mL.

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491