


CHAMOT

重组人骨形态发生蛋白-4 /Recombinant BMP-4,
Human, Animal-Free

CM065-5HP



CM065-20HP



CM065-100HP



CM065-500HP



CM065-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1

产品简介

2

产品使用

3

产品储存/运输

4

实验数据展示

重组人骨形态发生蛋白-4 /Recombinant BMP-4, Human, Animal-Free

产品编号	CM065-5HP	CM065-20HP	CM065-100HP	CM065-500HP	CM065-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述	Bone Morphogenetic Protein-4 (BMP-4) is an extracellular multifunctional signaling cytokine that is also a member of the TGF- β family. BMP-4 can bind with the TGF- β receptor and trigger SMAD protein signal transduction. BMP-4 has a significant expression in the prostate, it plays a vital role in the development of mesoderm into bones and the development of the epidermis and Dorsal-Ventral axis.
别称	bone morphogenetic protein 4, BMP-2B, DVR4, BMP4
蛋白编码	P12644
分子量	The protein has a calculated MW of 12.88 kDa. The protein migrates as 12 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>95% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce alkaline phosphatase production by ATDC5 cells. The ED ₅₀ for this effect is <0.58 ng/mL.
内毒素检测	<0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列	MKKKNKNRRHSLYVDFSDVGWNDWIVAPPGYQAFYCHGDCPFPLADHLNSTNHAIVQTLVNSVNNSIPKACCVPTELSAISMLYLDEYDKVVLKNYQEMVVEGCGCR with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 1×PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.
Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ($\geq 20 \mu\text{l}$).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant human BMP-4

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491