

CHAMOT

重组人成纤维细胞生长因子-17 /Recombinant FGF-17, Human, Animal-Free

CM106-5HP



CM106-20HP



CM106-100HP



CM106-500HP



CM106-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1

产品简介

2

产品使用

3

产品储存/运输

4

实验数据展示

重组人成纤维细胞生长因子-17 /Recombinant FGF-17, Human, Animal-Free

产品编号	CM106-5HP	CM106-20HP	CM106-100HP	CM106-500HP	CM106-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述	Fibroblast Growth Factors-17 (FGF-17) is a 23.7 kDa member of the fibroblast Growth Factors with 205 amino acid residues. FGF-17 is mainly expressed from brain, choroid plexus. FGF-17 involved embryonic development and cell proliferation. Required for normal brain development.
别称	fibroblast growth factor 17, FGFH, FGF17
蛋白编码	O60258
分子量	The protein has a calculated MW of 23.32 kDa. The protein migrates as 24 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce 3T3 cells proliferation. The ED ₅₀ for this effect is <5 ng/mL. The specific activity of recombinant human FGF-17 is > 2 x 10 ⁵ IU/mg.
内毒素检测	<0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列	TQGENHPSPNFNQYVRDQGAMTDQLSRRQIREYQLYSRTSGKHVQVTGRRISATAEDGNKF AKLIVETDTFGSRVRIKGAESEKYICMNKRGKLGKPSGKSKDCVFTIEVLENNYTAFQNARHEG WFMAFTRQGRPRQASRSRQNQREAHFIKRLYQGQLPFPNHAEKQKQFEFVGSAPTRRTKRT RPQPLT with polyhistidine tag at the N-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.
Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ($\geq 20 \mu\text{l}$).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant human FGF-17

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491