

# CHAMOT

重组人成纤维细胞生长因子-20 /Recombinant FGF-20, Human, Animal-Free

CM109-5HP



CM109-20HP



CM109-100HP



CM109-500HP



CM109-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

# CONTENT

1

产品简介

2

产品使用

3

产品储存/运输

4

实验数据展示

## 重组人成纤维细胞生长因子-20 /Recombinant FGF-20, Human, Animal-Free

产品编号	CM109-5HP	CM109-20HP	CM109-100HP	CM109-500HP	CM109-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

### 产品简介

背景描述	Fibroblast Growth Factors-20 (FGF-20) is a 23.5 kDa member of the fibroblast Growth Factors with 211 amino acid residues. FGF-20 is mainly expressed from microglial cells. FGF-20 can regulate central nervous development and function, regulate the survival of dopaminergic neurons via FGF/FGFR-dependent signaling pathway.
别称	fibroblast growth factor 20, FGFK, FGF20
蛋白编码	Q9NP95
分子量	The protein has a calculated MW of 24.24 kDa. The protein migrates as 24 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce 3T3 cells proliferation. The ED <sub>50</sub> for this effect is 1.3-3.2 ng/mL. The specific activity of recombinant human FGF-20 is > 2 x 10 <sup>5</sup> IU/mg.
内毒素检测	<0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列	MPLAEVGGFLGGLEGLGQQVGSHFLPPAGERPPLLGERRSAAERSARGGPGAAQLAHLHGILRRRQLYCRTGFHLQILPDGSVQGTRQDHSLFGILEFISAVGLVSIRGVDSGLYLGMDKGELYGSEKLTSECIFREQFEENWYNTYSSNIYKHGDTGRRYFVALNKDGTPRDGARSKRHQKFTHFLPRPVDPERVPELYKDLLMYT with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

### 产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

## 2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.  
**Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.**
- Store at 2°C to 8°C for up to 1 week.

## 3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ( $\geq 20 \mu\text{l}$ ).
- Store at -20°C or -80°C for 3 to 6 months.

**Avoid repeated freeze-thaw cycles.**

## 产品储存/运输

**Lyophilized Protein:** Store at -20°C for 1 year.


**储存** **Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

**Extended Storage:** Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

## 运输

Blue Ice

## 实验数据展示



SDS-PAGE analysis of recombinant human FGF-20

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. [www.chamot-bio.com](http://www.chamot-bio.com)

Tel: 021-51880030 Mail: [info@chamot-bio.com](mailto:info@chamot-bio.com) QQ: 864920491