

CHAMOT

重组人成纤维细胞生长因子-8a /Recombinant FGF-8a, Human, Animal-Free

CM096-5HP



CM096-20HP



CM096-100HP



CM096-500HP



CM096-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1

产品简介

2

产品使用

3

产品储存/运输

4

实验数据展示

重组人成纤维细胞生长因子-8a /Recombinant FGF-8a, Human, Animal-Free

产品编号	CM096-5HP	CM096-20HP	CM096-100HP	CM096-500HP	CM096-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述	Fibroblast Growth Factors-8a (FGF-8a) is a 23.5 kDa member of the fibroblast Growth Factors with 204 amino acid residues. FGF-8 is an important role in the regulation of embryonic development, cell proliferation, cell differentiation and cell migration. In embryogenesis function, FGF8 is necessary in limb, brain, eye and ear formation.
别称	fibroblast growth factor 8a, AIGF, HBGF-8, FGF8
蛋白编码	P55075
分子量	The protein has a calculated MW of 22.14 kDa. The protein migrates as 23 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce 3T3 cells proliferation. The ED ₅₀ for this effect is <2 µg/mL.
内毒素检测	<0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列	MQHVREQSLVTDQLSRRLIRTYQLYSRTSGKHVQVLANKRINAMAEDGDPFAKLIVETDTFGS RVRVRGAETGLYICMNKKGKLIAKSNGKGKDCVFTEIVLENNYTALQNAKYEGWYMAFTRKG RPRKGSKTRQHQREVHFMKRLPRGHHTEQSLRFEFNYPPFTRSLRGSQRTWAPEPR with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 50 mM Tris, 1 M NaCl, pH 8.5. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
 - Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.
- Do Not Vortex!** Vigorous shaking may impair the biological activity of the protein.
- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ($\geq 20 \mu\text{l}$).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant human FGF-

8a

Human FGF-8a, His Tag, E. coli induced 3T3 cell proliferation, with

the ED₅₀ at 1.93 µg/mL.

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491