

# CHAMOT

重组人高迁移率族蛋白B2 /Recombinant HMGB2,  
Human, Animal-Free

CM084-5HP



CM084-20HP



CM084-100HP



CM084-500HP



CM084-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

# CONTENTS

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

## 重组人高迁移率族蛋白B2 /Recombinant HMGB2, Human, Animal-Free

产品编号	CM084-5HP	CM084-20HP	CM084-100HP	CM084-500HP	CM084-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

### 产品简介

背景描述	High mobility group protein B2, also known as HMGB2, is a member of the non-histone chromosomal high-mobility group protein family. High mobility group protein B2 is expressed as 25 kDa containing 209 amino acid residues. The function of High mobility group protein B2 is involved in transcription, chromatin remodeling and V(D)J recombination. In vitro studies have demonstrated that this protein is able to efficiently bend DNA and form DNA circles.
别称	high mobility group box 2, HMG2
蛋白编码	P26583
分子量	The protein has a calculated MW of 24.84 kDa. The protein migrates as 34 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Testing in process
内毒素检测	<0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列	MGKGDPNKPRGKMSSYAFFVQTCREEHKKKHPDSSVNFAEFSKKCSERWKTMSAKEKSKFED MAKSDKARYDREMKNVPPKGDKKGKKDPNAPRPPSAFFLFCSEHRPKIKSEHPGLSIGDT AKKLGEMWSEQSAKDKQPYEQKAAKLKEKYEKDIAAYRAKGKSEAGKKGPGRPTGSKKNEP EDEEEEEEEDEDEEEDEEE with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

## 产品使用

### 1. Before opening, centrifuge at 3000 rpm for 5 mins.

### 2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

**Do Not Vortex!** Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

### 3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ( $\geq 20 \mu\text{l}$ ).
- Store at -20°C or -80°C for 3 to 6 months.

**Avoid repeated freeze-thaw cycles.**

## 产品储存/运输

**Lyophilized Protein:** Store at -20°C for 1 year.

### 储存


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

**Extended Storage:** Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

### 运输

Blue Ice

## 实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. [www.chamot-bio.com](http://www.chamot-bio.com)

Tel: 021-51880030 Mail: [info@chamot-bio.com](mailto:info@chamot-bio.com) QQ: 864920491