

CHAMOT

**重组人干扰素alpha 1a /Recombinant IFN alpha 1a,
Human, Animal-Free**

CM079-5HP

CM079-20HP

CM079-100HP

CM079-500HP

CM079-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组人干扰素alpha 1a /Recombinant IFN alpha 1a, Human, Animal-Free

产品编号	CM079-5HP	CM079-20HP	CM079-100HP	CM079-500HP	CM079-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述 Interferon-alpha 1a (IFN-alpha 1a) is a leukocyte interferon, which is a variant of Interferon-alpha. IFN-alpha 1a is a 19.5kDa protein containing 167 amino acid residues. It could bind the interferon receptors that activates the signal transduction of immune responses through the Jak/STAT pathway in leukocytes and lymphoblastoid cells.

别称 interferon alpha 1a, Leukocyte Interferon, IFNA2, B cell Interferon, Type I Interferon

蛋白编码 P01562

分子量 The protein has a calculated MW of 20.19 kDa. The protein migrates as 19 kDa under reducing condition (SDS-PAGE analysis).

表达系统 Escherichia coli

纯度 >95% as determined by SDS-PAGE.

生物活性 Measure by its ability to inhibit IL-8 secretion in human PBMCs in the presence of LPS. The ED₅₀ for this effect is <1.12 µg /mL.

内毒素检测 <0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列 CDLPETHSLDNRRTLMLLAQMSRISPSSCLMDRHDFGFPQEEFDGNQFQKAPASVLHELIQQ IFNLFTTKDSSAAWDEDLLDKFCTELYQQLNDLEACVMQEERVGETPLMNADSILAVKKYFRRI TLYLTEKKYSPCAWEVVRAEIMRSLSLSTNLQERLRRKE with polyhistidine tag at the N-terminus.

产品形式 The protein was lyophilized from a 0.2 µm filtered solution containing 500 mM NaCl PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

产品应用 Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/mL.

- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491