

# CHAMOT

**重组人白细胞介素11 /Recombinant IL-11, Human,  
Animal-Free**

CM013-5HP

CM013-20HP

CM013-100HP

CM013-500HP

CM013-1000HP



CHAMOT

乔默®生物

*Specialize In Cytokines*



# CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

## 重组人白细胞介素11 /Recombinant IL-11, Human, Animal-Free

产品编号	CM013-5HP	CM013-20HP	CM013-100HP	CM013-500HP	CM013-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

### 产品简介

**背景描述** Interleukin 11 (IL-11) is a cytokine, encoded by the gp130 family genes. IL-11 initially thought to be important for hematopoiesis, notably for megakaryocyte maturation, and it facilitate the formation of higher order structures involving dimers of gp130: IL-11:IL-11 RA complexes.

**别称** interleukin 11, AGIF (Adipogenesis Inhibitory Factor), IL11

**蛋白编码** P20809

**分子量** The protein has a calculated MW of 19.95 kDa. The protein migrates as 24 kDa under reducing condition (SDS-PAGE analysis).

**表达系统** Escherichia coli

**纯度** >98% as determined by SDS-PAGE.

**生物活性** Measure by its ability to induce T11 cells proliferation. The ED<sub>50</sub> for this effect is <0.2 ng/mL. The specific activity of recombinant human IL-11 is approximately >1 x 10<sup>7</sup> IU/ mg.

**内毒素检测** <0.1 EU per 1 µg of the protein by the LAL method.

**蛋白序列** PGPPPGPPRVSPDPRAELDSTVLLTRSLADTRQLAAQLRDKFPADGDHNLDSLPTLAMSAGALGALQLPGVLTRLRADLLSYLRHVQWLRRAGGSSKLTLEPELGTLQARLDRLRLRLQLLMSRLALPQPPDPAPPLAPPSSAWGGIRAAHAILGGLHLTDWAVRGLLLKTRL with polyhistidine tag at the N-terminus.

**产品形式** The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.

**产品应用** Cell Culture

## 产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

### 2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

### 3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/mL.

- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

## 产品储存/运输

**Lyophilized Protein:** Store at -20°C for 1 year.

### 储存


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

**Extended Storage:** Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

### 运输

Blue Ice

## 实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. [www.chamot-bio.com](http://www.chamot-bio.com)

Tel: 021-51880030 Mail: [info@chamot-bio.com](mailto:info@chamot-bio.com) QQ: 864920491