

CHAMOT

**重组人白细胞介素13 /Recombinant IL-13, Human,
Animal-Free**

CM015-5HP

CM015-20HP

CM015-100HP

CM015-500HP

CM015-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组人白细胞介素13 /Recombinant IL-13, Human, Animal-Free

产品编号	CM015-5HP	CM015-20HP	CM015-100HP	CM015-500HP	CM015-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述

Interleukin 13 (IL-13) with a molecular mass of 10 kDa cytokine, it secreted by T helper type 2 (Th2) cells, CD4 cells, natural killer T cell, mast cells, basophils, eosinophils and nuocytes. It is homologous to IL-4 and shares many of its biologic activities on mononuclear phagocytic cells, endothelial cells, epithelial cells, and B cells.

别称

interleukin 13, NC30, IL13

蛋白编码

P35225

分子量

The protein has a calculated MW of 13.28 kDa. The protein migrates as 15 kDa under reducing condition (SDS-PAGE analysis).

表达系统

Escherichia coli

纯度

>95% as determined by SDS-PAGE.

生物活性

Measure by its ability to induce TF-1 cells proliferation. The ED₅₀ for this effect is <0.8 ng/mL. The specific activity of recombinant human IL-13 is approximately >1 x10⁶ IU/ mg.

内毒素检测

<0.01 EU per 1 µg of the protein by the LAL method.

蛋白序列

MGPVPPSTALRELIEELVNITQNQKAPLCNGSMVWSINLTAGMYCAALES LINVSGCSAIEKTQ RMLSGFCPHKVSAGQFSSLHVRDTKIEVAQFVKDLLHLKKLFREGRFN with polyhistidine tag at the C-terminus

产品形式

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.

产品应用

Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.

- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491