


# CHAMOT

重组人白细胞介素16 /Recombinant IL-16 (aa 2-130), Human, Animal-Free

CM017-5HP



CM017-20HP



CM017-100HP



CM017-500HP



CM017-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

# CONTENTS

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

## 重组人白细胞介素16 /Recombinant IL-16 (aa 2-130), Human, Animal-Free

产品编号	CM017-5HP	CM017-20HP	CM017-100HP	CM017-500HP	CM017-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

### 产品简介

背景描述	Interleukin 16(IL-16) predicts a molecular mass of 13.5 kDa, encoded by this gene is a pleiotropic cytokine that functions as a chemoattractant, a modulator of T cell activation, and an inhibitor of HIV replication, and signaling process of this cytokine is mediated by CD4.
别称	interleukin 16, LCF (Lymphocyte Chemoattractant Factor), IL16
蛋白编码	Q14005
分子量	The protein has a calculated MW of 14.12 kDa. The protein migrates as 18 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>95% as determined by SDS-PAGE.
生物活性	Testing in process
内毒素检测	<0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列	MPDLNSSTDAAAASASAASDVSVESTAVCTVTLEKMSAGLGFSLGGKGS LHGDKPLTINRIF KGAASEQSETVQPGDEILQLGGTAMQGLTRFEAWNIIKALPDGPVTIVIRRKSLQS KETTAAGD S with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

### 产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

## 2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.  
**Do Not Vortex!** Vigorous shaking may impair the biological activity of the protein.
- Store at 2°C to 8°C for up to 1 week.

## 3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ( $\geq 20 \mu\text{l}$ ).
- Store at -20°C or -80°C for 3 to 6 months.

**Avoid repeated freeze-thaw cycles.**

## 产品储存/运输

**Lyophilized Protein:** Store at -20°C for 1 year.

### 储存


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

**Extended Storage:** Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

### 运输

Blue Ice

## 实验数据展示



SDS-PAGE analysis of recombinant human IL-16

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. [www.chamot-bio.com](http://www.chamot-bio.com)

Tel: 021-51880030 Mail: [info@chamot-bio.com](mailto:info@chamot-bio.com) QQ: 864920491