


CHAMOT

重组人白细胞介素9 /Recombinant IL-9, Human, Animal-Free

CM011-5HP



CM011-20HP




CM011-100HP



CM011-500HP



CM011-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENTS

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组人白细胞介素9 /Recombinant IL-9, Human, Animal-Free

产品编号	CM011-5HP	CM011-20HP	CM011-100HP	CM011-500HP	CM011-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述	Interleukin 9 (IL-9) is a pleiotropic cytokine that had pleiotropic functions in the immune system, has a molecular mass of 14.5 kDa. The major source of IL-9 is T lymphocytes. It is secreted by CD4+ helper cells that acts as a regulator of a variety of hematopoietic cells.
别称	interleukin 9, p40 cytokine, T-cell growth factor p40, HP40, IL9
蛋白编码	P15248
分子量	The protein has a calculated MW of 14.93 kDa. The protein migrates as 13-17 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce proliferation in MO7e cells. The ED ₅₀ for this effect is <0.25 ng/mL. The specific activity of recombinant human IL-9 is approximately >5 x10 ⁶ IU/ mg.
内毒素检测	<0.01 EU per 1 µg of the protein by the LAL method.
蛋白序列	QGCPTLAGILDINFLINKMQEDPASKCHCSANVTSCCLGIPSDNCTRPCFSERLSQMTNTTM QTRYPLIFSRVKKSVVLKNNKCPYFSCEQPCNQTTAGNALTFLKSLLEIFQKEKMRGMRGKI with polyhistidine tag at the N-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.
Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.
- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ($\geq 20 \mu\text{l}$).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant human IL-9

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491