

CHAMOT

重组人血小板生成素 /Recombinant TPO, Human, Animal-Free

CM124-5HP



CM124-20HP



CM124-100HP



CM124-500HP



CM124-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENTS

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组人血小板生成素 /Recombinant TPO, Human, Animal-Free

产品编号	CM124-5HP	CM124-20HP	CM124-100HP	CM124-500HP	CM124-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述	Thrombopoietin (TPO) is a glycoprotein that produced by the liver, kidney, marrow stroma and several other tissues. The TPO level in the blood is mostly negatively correlated with the abundance of platelets and bone marrow megakaryocytes, although multiple states of inflammation or infection, liver failure, and hematological disturbances are associated with unexpectedly high or low circulating levels of the hormone.
别称	thrombopoietin, Megakaryocyte colony-stimulating factor, c-MPL Ligand, MGDF, THPO
蛋白编码	P40225
分子量	The protein has a calculated MW of 19.6 kDa. The protein migrates as 20 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>95% as determined by SDS-PAGE.
生物活性	Testing in process
内毒素检测	< 0.01 EU per 1 µg of the protein by the LAL method.
蛋白序列	SPAPPACDLRVLSKLLRDSHVLHSRLSQCPEVHPLPTPVLLPAVDFSLGEWKTQMEETKAQDIL GAVTLLLEGVMAARGQLGPTCLSSLLGQLSGQVRLLL GALQSLLGTLPPQGRTTAHKDPNAIF LSFQHLLRGKVRFLMLVGGSTLCVRRAPPTTAVPSRTSLVLTNEL with polyhistidine tag at the N-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ($\geq 20 \mu\text{l}$).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491