


# CHAMOT

重组人TNF相关的凋亡弱诱导剂 / Recombinant  
TWEAK, Human, Animal-Free

CM062-5HP



CM062-20HP



CM062-100HP



CM062-500HP



CM062-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

# CONTENTS

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

## 重组人TNF相关的凋亡弱诱导剂 / Recombinant TWEAK, Human, Animal-Free

产品编号	CM062-5HP	CM062-20HP	CM062-100HP	CM062-500HP	CM062-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

### 产品简介

**背景描述** TNF-related weak inducer of apoptosis (TWEAK) is a type II membrane protein of the tumor necrosis factor (TNF) family members, high levels expression in human brain, muscle, heart, spleen, and thymus. TWEAK is a 17.3 kDa protein containing 129 residues. TWEAK induce apoptosis through cognate receptor Fn14 activate caspase-8 and caspase-3, in HSC3 cells KATO-III gastric adenocarcinoma cells and IFN- $\gamma$ -treated HT-29. TWEAK stimulate inflammatory cytokines in HT-29, A375 cells, WI-38 fibroblast sand astrocytes.

别称	TNF-related weak inducer of apoptosis, TNFSF12, DR3LG, Apo3 Ligand
蛋白编码	O43508
分子量	The protein has a calculated MW of 17.88 kDa. The protein migrates as 18 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce proliferation in HUVEC cells. The ED <sub>50</sub> for this effect is <6 ng/mL.
内毒素检测	<0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列	MKGRKTRARRAIAAHYEVHPRPGDGAQAGVDGTSGWEEARINSSPLRYNRQIGEFIVTRAGLYYLYCQVHFDEGKAVYLKLDLLVDGVLAIRCLEEFSATAASSLGPQLRLCQVSGLLALRPGSSLRIRTLPW AHLKAAPFLTYFGLFQVH with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

## 产品使用

### 1. Before opening, centrifuge at 3000 rpm for 5 mins.

### 2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.  
*Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.*
- Store at 2°C to 8°C for up to 1 week.

### 3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ( $\geq 20 \mu\text{l}$ ).
- Store at -20°C or -80°C for 3 to 6 months.

*Avoid repeated freeze-thaw cycles.*

## 产品储存/运输

**Lyophilized Protein:** Store at -20°C for 1 year.

### 储存


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

**Extended Storage:** Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

### 运输

Blue Ice

## 实验数据展示



SDS-PAGE analysis of recombinant human  
TWEAK

Human TWEAK, His Tag, E. coli induced HUVEC cell proliferation,  
with the ED<sub>50</sub> at 5.84 ng/mL.

For Research Use or Further Manufacturing Only  
Chamot Biotechnology(Shanghai) Co., Ltd. [www.chamot-bio.com](http://www.chamot-bio.com)  
Tel: 021-51880030 Mail: [info@chamot-bio.com](mailto:info@chamot-bio.com) QQ: 864920491