

CHAMOT

重组小鼠白细胞介素1 beta /Recombinant IL-1 beta,
Mouse, Animal-Free

CM002-5MP



CM002-20MP



CM002-100MP



CM002-500MP



CM002-1000MP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENTS

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

5 文献引用

重组小鼠白细胞介素1 beta /Recombinant IL-1 beta, Mouse, Animal-Free

产品编号	CM002-5MP	CM002-20MP	CM002-100MP	CM002-500MP	CM002-1000MP
规格	5 μ g	20 μ g	100 μ g	500 μ g	1 mg

产品简介

背景描述	Interleukin-1 beta (IL-1 β) is a major cytokine expressed in macrophage, NK cells, monocytes, and neutrophils. It also plays fundamental role in inflammatory response, including monocyte activation, which is essential for the host defense and pathogen and pathogen resistance. Pro-IL-1 β is cleaved by cytosolic caspase 1 to form mature IL-1 β .
别称	interleukin 1 beta, Catabolin, Lymphocyte-Activating Factor (LAF), Endogenous Pyrogen (EP), Leukocyte Endogenous Mediator (LEM), Mononuclear Cell Factor (MCF), IL1, IL1B
蛋白编码	P10749
分子量	The protein has a calculated MW of 18.3 kDa. The protein migrates as 17 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce D10.G4.1 cells proliferation. The ED ₅₀ for this effect is <8 pg/mL. The specific activity of recombinant mouse IL-1 beta is approximately >1.2x 10 ⁸ IU/mg.
内毒素检测	<0.1 EU per 1 μ g of the protein by the LAL method.
蛋白序列	MVPIRQLHYRLRDEQQKSLVLSDPYELKALHLNGQNIQQVIFSMSFVQGEPSNDKIPVALGLKGKNLYLSCVMKDGTPTLQLESVDPKQYPKKMEKRFVFNKIEVKSKEFESAEPNWWYISTSQAEHKPVFLGNNSGQDIIDFTMESVSS with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 μ m filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ($\geq 20 \mu\text{l}$).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



文献引用

- RNF115 Inhibits the Post-ER Trafficking of TLRs and TLRs-Mediated Immune Responses by Catalyzing K11-Linked Ubiquitination of RAB1A and RAB13. *Adv Sci (Weinh)*. 2022 Mar 28;e2105391. (IF 15.444) (Mouse IL-1 β)

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491