

CHAMOT

**重组小鼠白细胞介素17F /Recombinant IL-17F,
Mouse, Animal-Free**

CM015-5MP



CM015-20MP



CM015-100MP



CM015-500MP



CM015-1000MP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENTS

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组小鼠白细胞介素17F /Recombinant IL-17F, Mouse, Animal-Free

| 产品编号 | CM015-5MP | CM015-20MP | CM015-100MP | CM015-500MP | CM015-1000MP |
|------|-----------|------------|-------------|-------------|--------------|
| 规格 | 5 µg | 20 µg | 100 µg | 500 µg | 1 mg |

产品简介

| | |
|--------------|--|
| 背景描述 | Interleukin 17F (IL-17F) predicts a molecular mass of 30.1 kDa, is a cytokine of innate and adaptive immune system involved in maintenance of tissue integrity and antimicrobial host defense against infection by inducing the expression of genes that encode other proinflammatory cytokines, such as tumor necrosis factor, interleukin 1, interleukin 6 and some members of the colony-stimulating factor family. |
| 别称 | interleukin 17F, C87042, IL17, IL17F |
| 蛋白编码 | Q7TNI7 |
| 分子量 | The protein has a calculated MW of 15.82 kDa. The protein migrates as 17-25 kDa under reducing condition (SDS-PAGE analysis). |
| 表达系统 | Escherichia coli |
| 纯度 | >98% as determined by SDS-PAGE. |
| 生物活性 | Measure by its ability to induce IL-6 secretion in 3T3 cells. The ED ₅₀ for this effect is <100 ng/mL. |
| 内毒素检测 | <0.1 EU per 1 µg of the protein by the LAL method. |
| 蛋白序列 | MRKNPKAGVPALQKAGNCPPLEDNTVRVDIRIFNQNQGISVPREFQNRSSSPWDYNITRDP HRFPSEIAEAQCRHSGCINAQGQEDSTMNSVAIQQEILVLRREPQGCSNSFRLEKMLLKVGCT CVKPIVHQAA with polyhistidine tag at the C-terminus. |
| 产品形式 | The protein was lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate, 0.2 M NaCl, pH 4.5. If you have any concerns or special requirements, please confirm with us. |
| 产品应用 | Cell Culture |

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ($\geq 20 \mu\text{l}$).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491