

CHAMOT

重组小鼠白细胞介素7 /Recombinant IL-7, Mouse, Animal-Free

CM008-5MP



CM008-20MP



CM008-100MP



CM008-500MP



CM008-1000MP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1

产品简介

2

产品使用

3

产品储存/运输

4

实验数据展示

重组小鼠白细胞介素7 /Recombinant IL-7, Mouse, Animal-Free

产品编号	CM008-5MP	CM008-20MP	CM008-100MP	CM008-500MP	CM008-1000MP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述	Interleukin-7 (IL-7) is a multipotent cytokine belonged to one of the members of IL-2 superfamily. IL-7 has diverse effects on the hematopoietic and immune regulations. IL-7 is a trophic factor that is necessary for both B cell and T cell proliferation and development. In addition, it presents potential antitumor effects in tumors such as glioma, melanoma, lymphoma, leukemia, prostate cancer, and glioblastoma.
别称	interleukin 7, Lymphopoietin 1(LP-1), pre-B-cell factor, hlb368, A630026I06Rik, IL7
蛋白编码	P10168
分子量	The protein has a calculated MW of 15.8 kDa. The protein migrates as 16 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Measured in a cell proliferation assay using PHA-activated human peripheral blood lymphocytes (PBMC). The ED ₅₀ for this effect is <0.2 ng/mL. The specific activity of recombinant mouse IL-7 is > 5 x 10 ⁶ IU/mg.
内毒素检测	<0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列	MECHIKDKEGKAYESVLMISIDEELKMTGTDSCPNNEPNFFRKHVCDDTKEAAFLNRAARKLKQFLKMNISEEFNVHLLTVSQGTQTLVNCTSKEEKNVKEQKKNDACFLKRLREIKTCWNKILKGSI with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots ($\geq 20 \mu\text{l}$).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491