CHAMOT

重组小鼠Noggin /Recombinant Noggin, Mouse, AF, HEK293

CM099-5MP

CM099-20MP

CM099-100MP

CM099-500MP

CM099-1000MP





トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



Product Datasheet

重组小鼠Noggin /Recombinant Noggin, Mouse, AF, HEK293

| 产品编号 | CM099-5MP | CM099-20MP | CM099-100MP | CM099-500MP | CM099- 1000MP |
|------|-----------|------------|-------------|-------------|------------------|
| 规格 | 5 μg | 20 μg | 100 μg | 500 μg | 1 mg |

产品简介

Noggin binds members of the transforming Growth Factors-beta (TGF beta) superfamily signaling proteins, such as bone morphogenetic protein-4 (BMP-4), which inactivates their activities. As a extracellular antagonist of BMP proteins, Noggin involves in the development of many body tissues, including nerve tissue, muscles, and bones. In addition, Noggin is able to inhibit chondrocyte differentiation through its interaction with GDE5.

| | through its interaction with GDF5. |
|-------|--|
| 别称 | Nog |
| 蛋白编码 | P97466 |
| 分子量 | The protein has a calculated MW of 49.14 kDa. The protein migrates as 50 kDa under reducing condition (SDS-PAGE analysis). |
| 表达系统 | HEK293 cell |
| 纯度 | >95% as determined by SDS-PAGE. |
| 生物活性 | Testing in process |
| 内毒素检测 | $<$ 0.1 EU per 1 μ g of the protein by the LAL method. |
| 蛋白序列 | A DNA sequence encoding Mouse Noggin(#P97466) (Met1-Cys232) was expressed with Human IgG1 Fc tag at the C-terminus. |
| 产品形式 | The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH |

7.4. If you have any concerns or special requirements, please confirm with us.

产品使用

产品应用

Cell Culture

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than $10~\mu g/ml$.
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存 Initial Reconstitution

Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.


Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant mouse Noggin

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491