

CHAMOT

**重组猪白细胞介素2 /Recombinant IL-2, Swine,
Animal-Free**

CM003-5SP

CM003-20SP

CM003-100SP

CM003-500SP

CM003-1000SP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组猪白细胞介素2 /Recombinant IL-2, Swine, Animal-Free

产品编号	CM003-5SP	CM003-20SP	CM003-100SP	CM003-500SP	CM003-1000SP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述 Interleukin-2 (IL-2) is a four bundle, immuno-modulatory cytokine which is expressed predominately by antigen-simulated CD4+ T cells, CD8+ T cells, natural killer (NK) cells, and dendritic cells (DC). IL-2 plays a fundamental role in promoting NK cell proliferation, cytotoxicity, and B cells differentiation. IL-2 modulate T cell differentiation programs, promoting naïve CD4+ T cell differentiation into T helper-1 (Th1) and T helper-2 (Th2) cells.

别称 interleukin 2, T-cell Growth Factor (TCGF), Aldesleukin, POIL2

蛋白编码 P26891

分子量 The protein has a calculated MW of 16.2 kDa. The protein migrates as 13 kDa under reducing condition (SDS-PAGE analysis).

表达系统 Escherichia coli

纯度 >98% as determined by SDS-PAGE.

生物活性 Measure by its ability to induce proliferation in CTLL2 cells. The ED₅₀ for this effect is <0.5 ng/mL.

内毒素检测 <0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列 MAPTSSSTKNTKKQLEPLLLDLQLLLKEVKNYENADLSRMLTFKFYMPKQATELKHLQCLVEEL KALEGVNLGQSKNSDSANIKESMNNINVTVLELKGSETSFKCEYDDETVTAVEFLNKWITFCQ SIYSTLT with polyhistidine tag at the C-terminus

产品形式 The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

产品应用 Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant swine IL-2

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491