# **CHAMOT**

## 重组猪白细胞介素6 /Recombinant IL-6, Swine, Animal-Free

CM005-5SP

CM005-20SP

CM005-100SP

CM005-500SP

CM005-1000SP





# トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



### **Product Datasheet**

### 重组猪白细胞介素6 /Recombinant IL-6, Swine, Animal-Free

产品编号	CM005-5SP	CM005-20SP	CM005-100SP	CM005-500SP	CM005-1000SP
规格	5 μg	20 μg	100 µg	500 μg	1 mg

_		
	ALA.	
_		
- 1 -		

背景描述

分子量

Interleukin-6 (IL-6) is a pleiotropic cytokine which plays fundamental role in the acute phase response, inflammation, bone metabolism, lymphocyte differentiation and cancer progression. Deregulation of IL-6 production was also found in several diseases, including rheumatoid arthritis, Alzheimer's disease, autoimmune deficiency disease and different types of cancer.

别称 interleukin 6, IL6

P26893 蛋白编码

The protein has a calculated MW of 21.9 kDa. The protein migrates as 23 kDa under

reducing condition (SDS-PAGE analysis).

表达系统 Escherichia coli

纯度 >98% as determined by SDS-PAGE.

Measure by its ability to induce proliferation in T1165.85.2.1 cells. The ED50 for this 生物活性

effect is <1.5 ng/mL.

内毒素检测 < 0.1 EU per 1 µg of the protein by the LAL method.

MGRLEEDAKGDATSDKMLFTSPDKTEELIKYILGKISAMRKEMCEKYEKCENSKEVLAENNLNL PKMAEKDGCFQSGFNQETCLMRITTGLVEFQIYLDYLQKEYESNKGNVEAVQISTKALIQTLRQ

KGKNPDKATTPNPTTNAGLLDKLQSQNEWMKNTKIILILRSLEDFLQFSLRAIRIM with

polyhistidine tag at the C-terminus.

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 产品形式

7.4. If you have any concerns or special requirements, please confirm with us.

产品应用 Cell Culture

蛋白序列

**1.** Before opening, centrifuge at 3000 rpm for 5 mins.

### 2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

### 3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than  $10~\mu g/ml$ .
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

### 产品储存/运输

**Lyophilized Protein:** Store at -20°C for 1 year.

储存 Initial Reconstitution

**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.


**Extended Storage:** Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输

Blue Ice

### 实验数据展示



SDS-PAGE analysis of recombinant swine IL-6

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491