CHAMOT

重组小鼠白细胞介素5 /Recombinant IL-5, Mouse, Animal-Free

CM006-5MP

CM006-20MP

CM006-100MP

CM006-500MP

CM006-1000MP





トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



Product Datasheet

重组小鼠白细胞介素5 /Recombinant IL-5, Mouse, Animal-Free

产品编号	CM006-5MP	CM006-20MP	CM006-100MP	CM006-500MP	CM006- 1000MP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

产品简介

背景描述	Interleukin 5 (IL-5) is a 45-60 kDa homodimeric cytokine that functions principally in the eosinophil arm of the Th2 response. It is a member of the helical bundle family of cytokines, along with IL-3, IL-4, and granulocyte macrophage-colony stimulating factor (GM-CSF).
别称	interleukin 5, EDF, BCDFII, TRF, IL5
蛋白编码	P04401
分子量	The protein has a calculated MW of 13.93 kDa. The protein migrates as 11-17 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce TF-1 cells proliferation. The ED $_{50}$ for this effect is <0.2 ng/mL. The specific activity of recombinant mouse IL-5 is > 5 x 10 6 IU/mg.
内毒素检测	$<$ 0.1 EU per 1 μ g of the protein by the LAL method.
蛋白序列	MEIPMSTVVKETLTQLSAHRALLTSNETMRLPVPTHKNHQLCIGEIFQGLDILKNQTVRGGTVE MLFQNLSLIKKYIDRQKEKCGEERRRTRQFLDYLQEFLGVMSTEWAMEG with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 μ m filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than $10~\mu g/ml$.
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存 Initial Reconstitution:


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输 Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant mouse IL-5

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491