

CHAMOT

Goat anti-Human IgG (Fc Fragment)-HRP

CM008-A2HS

CM008-0.1A2H

CM008-0.5A2H

CM008-1A2H

CM008-5A2H

CM008-10A2H



CHAMOT

乔默®生物

Specialize In Cytokines



CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

Goat anti-Human IgG (Fc Fragment)-HRP

产品编号	CM008-A2HS	CM008-0.1A2H	CM008-0.5A2H	CM008-1A2H	CM008-5A2H	CM008-10A2H
规格	10 μL	100 μL	500 μL	1 mL	5 mL	10 mL

产品简介

产品描述	Goat anti-Human IgG, Fc Fragment is purified from goat antiserum by immunoaffinity chromatography and conjugated with high-activity HRP. Minimal cross-reactivity with other isotypes or species.
产品形式	Liquid
产品应用	ELISA,IHC,WB
反应性	Human
克隆性	Polyclonal
免疫原	IgG, Fc Fragment
浓度	Lot-dependent, please refer to product label.
标记	HRP
产品组成	Phosphate Buffered Saline containing 1% BSA, 0.03% ProClin 300, pH 7.4.
亚型	IgG, Fc Fragment

产品使用

Recommended dilution factor:
ELISA: 1:5000-25000
IHC: 1:500-2500
WB:1:5000-25000

Note: Working dilution for specific application should be determined by the investigator to obtain the best conditions.

产品储存/运输

储存

This product is stable after storage at: 2-8°C for 2 weeks under sterile conditions from date of receipt. -20°C or -80°C for 12 months under sterile conditions from date of receipt. Avoid repeated freeze/thaw cycles and protect from light. Suggestion: Divide antibody into several vials. Keep only vials for usage at 2-8°C.

运输

The product is shipped with polar packs. Upon receipt, store it immediately at -20°C or lower for long term storage.

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491