

CHAMOT

**重组猪肿瘤坏死因子alpha /Recombinant TNF
alpha, Swine, Animal-Free**

CM010-5SP

CM010-20SP

CM010-100SP

CM010-500SP

CM010-1000SP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组猪肿瘤坏死因子alpha /Recombinant TNF alpha, Swine, Animal-Free

产品编号	CM010-5SP	CM010-20SP	CM010-100SP	CM010-500SP	CM010-1000SP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述

Tumor necrosis factor alpha (TNF alpha) stimulates the acute phase of the immune response. In response to a pathogen, TNF alpha is one of the first to be released and can apply its effects in many organs. TNF alpha stimulates the release of corticotropic releasing hormone, suppresses appetite, and induces fever, in the hypothalamus. TNF increase vasodilation and loss of vascular permeability, it helps recruit lymphocyte, neutrophil, and monocyte to the inflammation site by regulating chemokine release.

别称

tumor necrosis factor alpha, TNFa, TNFSF2

蛋白编码

P23563

分子量

The protein has a calculated MW of 18.1 kDa. The protein migrates as 13 kDa under reducing condition (SDS-PAGE analysis).

表达系统

Escherichia coli

纯度

>98% as determined by SDS-PAGE.

生物活性

Measure by its ability to induce cytotoxicity in PK15 cells in the presence of the actinomycin D. The ED₅₀ for this effect is <15 pg/mL.

内毒素检测

<0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列

MLRSSSQTS DKPVAHVVANVKAEGQLQWQSGYANALLANGVVKLKDNLVPTDGLYLIYSQ
VLFRRGQGCPSTNVFLTHTISRIVASYQTKVNULLSAIKSPCQRETPEGAEAKPWYEPIYLGGVFQLE
KDDRLSAEINLPDYLDFAESGQVYFGIIAL with polyhistidine tag at the C-terminus.

产品形式

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

产品应用

Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2. Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3. Extended Storage

• After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.

- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491