

CHAMOT

**重组猪骨形态发生蛋白-4 /Recombinant BMP-4,
Swine, Animal-Free**

CM011-5SP

CM011-20SP

CM011-100SP

CM011-500SP

CM011-1000SP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组猪骨形态发生蛋白-4 /Recombinant BMP-4, Swine, Animal-Free

产品编号	CM011-5SP	CM011-20SP	CM011-100SP	CM011-500SP	CM011-1000SP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述

Bone Morphogenetic Protein-4 (BMP-4) predicts a molecular mass of 13 kDa, is a vital regulatory molecule that functions throughout human development in mesoderm induction, tooth development, limb formation, bone induction, and fracture repair and is overexpressed in patients who have fibrodysplasia ossificans progressiva. BMP-4 is a critical signaling molecule required for the early differentiation of the embryo and establishing of a dorsal-ventral axis. BMP-4 is secreted from the dorsal portion of the notochord, and it acts in concert with sonic hedgehog to establish a dorsal-ventral axis for the differentiation of later structures.

别称

bone morphogenetic protein 4, BMP4

蛋白编码

A7LJT9

分子量

The protein has a calculated MW of 21.65 kDa. The protein migrates as 17-25 kDa under reducing condition (SDS-PAGE analysis).

表达系统

Escherichia coli

纯度

>98% as determined by SDS-PAGE.

生物活性

Testing in process

内毒素检测

<0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列

MREKQPNYGLAIEVTHLHQTRTHQGQHVRISRSLPQGS GDWAQLRPLLVTFGHDGRGHALT
RRRRAKRSPKHHHPQRARKKNKNCRRHS LYVDFSDVGWNDWIVAPPGYQAFYCHGDCPFPL
ADHLNSTNHAIVQTLVNSVNSSIPKACCVPTELSAISMLYLDEYDKVVLKKNYQEMVVEGCGC
with polyhistidine tag at the C-terminus.

产品形式

The protein was lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate, 0.2 M NaCl, pH 3.5. If you have any concerns or special requirements, please confirm with us.

产品应用

Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.

- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491