# **CHAMOT**

重组小鼠白细胞介素24 /Recombinant IL-24, Mouse, Animal-Free


CM020-5MP

CM020-20MP

CM020-100MP

CM020-500MP

CM020-1000MP





# トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



### **Product Datasheet**

## 重组小鼠白细胞介素24 /Recombinant IL-24, Mouse, Animal-Free

产品编号	CM020-5MP	CM020-20MP	CM020-100MP	CM020-500MP	CM020- 1000MP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

### 产品简介

背景描述	Interleukin 24 (IL-24) predicts a molecular mass of 18.2 kDa, is a cytokine belonging to the IL-10 family of cytokines that signals through IL-20R1 /IL-20R2 and IL-22R1 /IL-20R2. Its normal physiological role is connected with wound healing, protection against diseases caused by bacteria.
别称	interleukin 24, MDA-7 (Melanoma Differentiation-Associated gene 7 protein), FISP, St16, IL24
蛋白编码	Q925S4
分子量	The protein has a calculated MW of 18.94 kDa. The protein migrates about 17 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>95% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce proliferation in BaF3 cells transfected with human IL-20 R beta. The ED $_{50}$ for this effect is <0.3 ng/mL.
内毒素检测	$<$ 0.1 EU per 1 $\mu$ g of the protein by the LAL method.
蛋白序列	MQEFRFGSCQVTGVVLPELWEAFWTVKNTVQTQDDITSIRLLKPQVLRNVSGAESCYLAHSLL KFYLNTVFKNYHSKIAKFKVLRSFSTLANNFIVIMSQLQPSKDNSMLPISESAHQRFLLFRRAFK QLDTEVALVKAFGEVDILLTWMQKFYHL with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 $\mu$ m filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

### 产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

### 2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

### 3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than  $10~\mu g/ml$ .
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

### 产品储存/运输

储存

Lyophilized Protein: Store at -20°C for 1 year.


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

**Extended Storage:** Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输 Blue Ice

### 实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491