

CHAMOT

**重组猪趋化因子CXCL10 /Recombinant CXCL10,
Swine, Animal-Free**

CM022-5SP

CM022-20SP

CM022-100SP

CM022-500SP

CM022-1000SP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组猪趋化因子CXCL10 /Recombinant CXCL10, Swine, Animal-Free

产品编号	CM022-5SP	CM022-20SP	CM022-100SP	CM022-500SP	CM022-1000SP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述

C-X-C motif chemokine 10 (CXCL10) also named Interferon gamma-induced protein 10 (IP-10), which is a chemokine of the intercrine alpha family. CXCL10 is a 9.1kDa protein containing 82 amino acid residues. CXCL10 is produced by the several cell types like monocytes and endothelial cells, which are responded for IFN γ . CXCL10 is a chemotaxis for T cells, NK cells and macrophages. CXCL10 also binds the CXCR3 that induce the the cell migration and activation like T cells and dendritic cells

别称

C-X-C motif chemokine 10, MCP-1

蛋白编码

NP_001008691 1

分子量

The protein has a calculated MW of 10.13 kDa. The protein migrates as 11-17 kDa under reducing condition (SDS-PAGE analysis).

表达系统

Escherichia coli

纯度

>98% as determined by SDS-PAGE.

生物活性

Measure by its ability to chemoattract BaF3 cells transfected with human CXCR2. The ED₅₀ for this effect is <4 ng/mL.

内毒素检测

<0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列

PLSRTVRCTCIKISDRPVNPRSLEKLEMIPASQSCPHVEIIATMKKNGEKRCLNPESKTIKNLLKAI SKERSKRSPRTQREA with polyhistidine tag at the N-terminus.

产品形式

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

产品应用

Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant swine CXCL10

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491