CHAMOT

重组小鼠白细胞介素28B /Recombinant IL-28B, Mouse, Animal-Free

CM025-5MP

CM025-20MP

CM025-100MP

CM025-500MP

CM025-1000MP





トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



Product Datasheet

重组小鼠白细胞介素28B / Recombinant IL-28B, Mouse, Animal-Free

| 产品编号 | CM025-5MP | CM025-20MP | CM025-100MP | CM025-500MP | CM025- 1000MP |
|------|-----------|------------|-------------|-------------|------------------|
| 规格 | 5 μg | 20 μg | 100 μg | 500 μg | 1 mg |

产品简介

产品应用

Cell Culture

| 背景描述 | Interleukin 28B (IL-28B) predicts a molecular mass of 19.7 kDa. IL-28 comes in two isoforms, IL-28A and IL-28B, and plays a role in immune defense against viruses, including the induction of an "antiviral state" by turning on Mx proteins, 2',5'-oligoadenylate synthetase as well as ISGF3G. |
|-------|---|
| 别称 | interleukin 28B, IFN-λ3, IFL, IFL-1, INF-alpha, INF-l, INF-lambda, IL28 |
| 蛋白编码 | Q8CGK6 |
| 分子量 | The protein has a calculated MW of 20.56 kDa. The protein migrates as 17-25 kDa under reducing condition (SDS-PAGE analysis). |
| 表达系统 | Escherichia coli |
| 纯度 | >98% as determined by SDS-PAGE. |
| 生物活性 | Measure by its ability to protect HepG2 cells infected with encephalomyocarditis (EMC) virus. The ED $_{50}$ for this effect is <20 ng/mL. |
| 内毒素检测 | $<$ 0.1 EU per 1 μ g of the protein by the LAL method. |
| 蛋白序列 | MDPVPRATRLPVEAKDCHIAQFKSLSPKELQAFKKAKGAIEKRLLEKDMRCSSHLISRAWDLKQ LQVQERPKALQAEVALTLKVWENINDSALTTILGQPLHTLSHIHSQLQTCTQLQATAEPKPPSRR LSRWLHRLQEAQSKETPGCLEDSVTSNLFQLLLRDLKCVASGDQCV with polyhistidine tag at the C-terminus. |
| 产品形式 | The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us. |
| | |

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than $10~\mu g/ml$.
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

储存

Lyophilized Protein: Store at -20°C for 1 year.


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输 Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491