CHAMOT

重组猪趋化因子CXCL11 /Recombinant CXCL11, Swine, Animal-Free

CM025-5SP

CM025-20SP

CM025-100SP

CM025-500SP

CM025-1000SP





トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



Product Datasheet

重组猪趋化因子CXCL11 / Recombinant CXCL11, Swine, Animal-Free

产品编号	CM025-5SP	CM025-20SP	CM025-100SP	CM025-500SP	CM025-1000SP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

产品简介

背景描述	C-X-C motif chemokine 11 (CXCL11) also named Interferon-gamma-inducible protein 9 (IP-9), which is a chemokine of the intercrine alpha family. CXCL11 is a 8.8kDa protein containing 79 amino acid residues. To CXCR3, CXCL11 has higher affinity than CXCL10 and CXCL9 which plays a role in immune activation. CXCL11 induces the activation of T cells which is also a chemotaxis for T cells. CXCL11 is produced in response for IFN Family.	
别称	C-X-C motif chemokine 11, MCP-1	
蛋白编码	NP_001121963 1	
分子量	The protein has a calculated MW of 9.72 kDa. The protein migrates about 11 kDa under reducing condition (SDS-PAGE analysis).	
表达系统	Escherichia coli	
纯度	>98% as determined by SDS-PAGE.	
生物活性	Testing in process	
内毒素检测	$<$ 0.1 EU per 1 μ g of the protein by the LAL method.	
蛋白序列	FPMFKAGRCLCIGPGVKAVKVADIEKVSIIHPSNNCDKTEVIVTLKAHKGRRCLNPKSKQANVI MKKVERMNFLRYQNV with polyhistidine tag at the N-terminus.	
产品形式	The protein was lyophilized from a 0.2 μ m filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.	
产品应用	Cell Culture	

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than $10~\mu g/ml$.
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存 Initial Reconstitution:


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输 Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant swine CXCL11

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491