CHAMOT

重组人白细胞介素25 /Recombinant IL-25, Human, Animal-Free


CM029-5HP

CM029-20HP

CM029-100HP

CM029-500HP

CM029-1000HP





トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



产品简介

Product Datasheet

重组人白细胞介素25 /Recombinant IL-25, Human, Animal-Free

产品编号	CM029-5HP	CM029-20HP	CM029-100HP	CM029-500HP	CM029-1000HP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

背景描述	Interleukin 25 (IL-25) is a cytokine with proinflammatory and anti-inflammatory effects, it predicts a molecular mass of 16.7 kDa. It can induce NF-κB activation, and stimulate the production of IL-8, which is the major chemotactic substance of neutrophils.
别称	interleukin 25, IL17E, IL25, IL-17E
蛋白编码	Q9H293
分子量	The protein has a calculated MW of 17.68 kDa. The protein migrates as 19 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce IL-8 secretion in human PBMCs. The ED $_{50}$ for this effect is <5 ng/mL. Measure by its ability to induce CXCL1 secretion in HT29 cells. The ED $_{50}$ for this effect is <1 ng/mL.
内毒素检测	$<$ 0.1 EU per 1 μ g of the protein by the LAL method.
	MYSHWPSCCPSKGQDTSEELLRWSTVPVPPLEPARPNRHPESCRASEDGPLNSRAISPWRYEL

DRDLNRLPQDLYHARCLCPHCVSLQTGSHMDPRGNSELLYHNQTVFYRRPCHGEKGTHKGY

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH

8.0. If you have any concerns or special requirements, please confirm with us.

CLERRLYRVSLACVCVRPRVMG with polyhistidine tag at the C-terminus.

产品使用

蛋白序列

产品形式

产品应用

Cell Culture

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than $10~\mu g/ml$.
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存 Initial Reconstitution


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输 Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant human IL-25

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491