

CHAMOT

**重组人白细胞介素32 alpha /Recombinant IL-32
alpha, Human, Animal-Free**

CM037-5HP

CM037-20HP

CM037-100HP

CM037-500HP

CM037-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组人白细胞介素32 alpha /Recombinant IL-32 alpha, Human, Animal-Free

产品编号	CM037-5HP	CM037-20HP	CM037-100HP	CM037-500HP	CM037-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述

Interleukin 32 alpha (IL-32 α) is an 18 kDa cytokine with 131 amino acid residues and one of the IL-2 splice variants. IL-32 α is primarily secreted from inflammatory cells, such as NK cells, T-cells, peripheral blood mononuclear cells (PBMCs), and monocytes. IL-32 alpha is a proinflammatory cytokine regulating IL-6 by interaction with protein kinase C(PKC). It also interacts with paxillin, integrin, and focal adhesion kinase 1 (FAK 1).

别称

interleukin 32 alpha, IL-32alpha, IL-32beta, IL-32delta, IL-32gamma, NK4, TAIF, TAIFa, TAIFb, TAIFc, TAIFd, IL32

蛋白编码

P24001-4

分子量

The protein has a calculated MW of 15.72 kDa. The protein migrates as 18 kDa under reducing condition (SDS-PAGE analysis).

表达系统

Escherichia coli

纯度

>98% as determined by SDS-PAGE.

生物活性

Measure by its ability to induce TNF alpha secretion in RAW264.7 cells. The ED₅₀ for this effect is <10 µg/mL.

内毒素检测

<0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列

MCFPKVLSDDMKKLKARMHQAIERFYDKMQNAESGRGQVMSSLAELEDDFKEGYLETVAAY YEEQHPELTPLEKERDGLRCRGNRSPVPDVEDPATEEPGESFCDKSYGAPRGDKEELTPQKCSE PQSSK with polyhistidine tag at the C-terminus .

产品形式

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.

产品应用

Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.

- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491