CHAMOT

重组小鼠白细胞介素38 /Recombinant IL-38, Mouse, Animal-Free

CM037-5MP

CM037-20MP

CM037-100MP

CM037-500MP

CM037-1000MP





トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



Product Datasheet

重组小鼠白细胞介素38 /Recombinant IL-38, Mouse, Animal-Free

产品编号	CM037-5MP	CM037-20MP	CM037-100MP	CM037-500MP	CM037- 1000MP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

产品简介

产品应用

Cell Culture

背景描述	Interleukin 38 (IL-38) is a 16.87 kDa cytokine with 152 amino acid residues and a member of the interleukin-1 (IL-1) family. IL-38 is primarily secreted from basal epithelia of the skin and B cells of the tonsil, spleen, and thymus. It plays essential roles in inflammation and host defense by mediating NF-kB, AP1, and JNK signaling pathways when IL-38 binds to the receptors like IL1RAPL1 and IL-36R. In addition, IL-38 participates in down-regulating cell proliferation, migration, and up-regulating IL-6 and IL-10.		
别称	interleukin 38, interleukin 1 family, member 10, IL1F10, IL38		
蛋白编码	Q8R459		
分子量	The protein has a calculated MW of 17.89 kDa. The protein migrates as 17-25 kDa under reducing condition (SDS-PAGE analysis).		
表达系统	Escherichia coli		
纯度	>98% as determined by SDS-PAGE.		
生物活性	Testing in process		
内毒素检测	$<$ 0.1 EU per 1 μ g of the protein by the LAL method.		
蛋白序列	MCSLPMARYYIIKDAHQKALYTRNGQLLLGDPDSDNYSPEKVCILPNRGLDRSKVPIFLGMQG GSCCLACVKTREGPLLQLEDVNIEDLYKGGEQTTRFTFFQRSLGSAFRLEAAACPGWFLCGPAE PQQPVQLTKESEPSTHTEFYFEMSR with polyhistidine tag at the C-terminus.		
产品形式	The protein was lyophilized from a 0.2 μ m filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.		

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than $10~\mu g/ml$.
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

储存

Lyophilized Protein: Store at -20°C for 1 year.


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输 Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491