

CHAMOT

**重组人白细胞介素33 /Recombinant IL-33, Human,
Animal-Free**

CM038-5HP

CM038-20HP

CM038-100HP

CM038-500HP

CM038-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组人白细胞介素33 /Recombinant IL-33, Human, Animal-Free

产品编号	CM038-5HP	CM038-20HP	CM038-100HP	CM038-500HP	CM038-1000HP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述

Interleukin 33 (IL-33) is an 18 kDa cytokine with 169 amino acid residues. Interleukin-33 is a member of the IL-1 cytokine family and is primarily expressed in endothelial cells, epithelial cells, fibroblasts, and mesenchymal cells. When IL-33 binds to its receptor, ST2L, it activates mast cells and innate lymphoid cells and subsequently generates IL-5 and IL-13, which play essential roles in allergic inflammation. IL-33 also acts as an alarm by alerting the immune system of cellular damage and plays a critical role in type-2 innate immunity.

别称

interleukin 33, IL-1 F11, NF-HEV, DVS27, IL33

蛋白编码

O95760

分子量

The protein has a calculated MW of 18.93 kDa. The protein migrates as 19 kDa under reducing condition (SDS-PAGE analysis).

表达系统

Escherichia coli

纯度

>98% as determined by SDS-PAGE.

生物活性

Measure by its ability to binding with recombinant ST2L (IL-33 receptor). The ED₅₀ for this effect is <54 ng/mL.

内毒素检测

<0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列

MSITGISPITEYLASLSTYNDQSITFALEDESYEIYVEDLKKDEKKDKVLLSYYESQHPSNESGDGV
DGKMLMVTLSPTKDFWLHANNKEHSVELHKCEKPLPDQAFFVLHNMHSNCVSFECKTDPG
VFIGVKDNHLALIKVDSSENLCTENILFKLSET with polyhistidine tag at the C-terminus.

产品形式

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

产品应用

Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.

- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491