

CHAMOT

**重组小鼠B细胞激活因子 /Recombinant BAFF,
Mouse, Animal-Free**

CM040-5MP

CM040-20MP

CM040-100MP

CM040-500MP

CM040-1000MP



CHAMOT

乔默®生物

Specialize In Cytokines



CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组小鼠B细胞激活因子 /Recombinant BAFF, Mouse, Animal-Free

产品编号	CM040-5MP	CM040-20MP	CM040-100MP	CM040-500MP	CM040-1000MP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述

BAFF also known as BLYS, TALL-1 and TNFSF13B, which belongs to tumor necrosis factor family. BAFF is a 31.2 kDa type II transmembrane protein containing 285 residues that predominantly produced by myeloid cells, furthermore mouse BAFF shares 72% sequence identity with human BAFF. BAFF has been demonstrated to activate the survival of B cells and the B cell response by binding to BAFFR/BR3. Additionally, BAFF also takes part in regulating B and T cell function via forming two ligands-two receptors pathway through sharing TNFRSF13B/TACI and TNFRSF17/BCMA receptors with APRIL.

别称

B-cell activating factor, tumor necrosis factor (ligand) superfamily, member 13b, Tnfsf13b, BAF, BL, BlyS, D8Ertd387, D8Ertd387e, TAL, TALL-1, TALL1, THANK, TNFSF20, Tnlg7a, zTNF, zTNF4

蛋白编码

Q9WU72

分子量

The protein has a calculated MW of 21.56 kDa. The protein migrates as 17-25 kDa under reducing condition (SDS-PAGE analysis).

表达系统

Escherichia coli

纯度

>98% as determined by SDS-PAGE.

生物活性

Measure by its ability to induce proliferation in mouse B cells. The ED₅₀ for this effect is <0.5 ng/mL. The specific activity of recombinant mouse BAFF is > 2 x 10⁶ IU/mg.

内毒素检测

<0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列

MAFQGPEETE QD V D L S A P P A P C L P G C R H S Q H D D N G M N L R N I I Q D C L Q L I A D S D T P T I R K G T Y T F V P W L L S F K R G N A L E E K E N K I V V R Q T G Y F F I Y S Q V L Y T D P I F A M G H V I Q R K K V H V F G D E L S L V T L F R C I Q N M P K T L P N N S C Y S A G I A R L E E G D E I Q L A I P R E N A Q I S R N G D D T F F G A L K L L with polyhistidine tag at the C-terminus.

产品形式

The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

产品应用

Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H_2O to a concentration of 100-200 $\mu\text{g}/\text{mL}$.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 $\mu\text{g}/\text{mL}$.

- Prepare aliquots ($\geq 20 \mu\text{l}$).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

储存

Lyophilized Protein: Store at -20°C for 1 year.


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant mouse BAFF

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491