CHAMOT

重组小鼠Fas配体 /Recombinant FasL,Mouse, Animal-Free


CM042-5MP

CM042-20MP

CM042-100MP

CM042-500MP

CM042-1000MP





トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

CHAMOT 乔默[®]生物

Product Datasheet

重组小鼠Fas配体 /Recombinant FasL, Mouse, Animal-Free

产品编号	CM042-5MP	CM042-20MP	CM042-100MP	CM042-500MP	CM042- 1000MP
规格	5 μg	20 μg	100 µg	500 μg	1 mg

产品简介

背景描述	Fas Ligand (FasL) is a 17.34 kDa tumor necrosis factor with 152 amino acid residues. FasL is mainly expressed from lymphoid tissue and secreted to blood. Binding to its receptor, TNFRSF6/FAS, leads to induce apoptotic signal into cells. Involved in cytotoxic T-cell-mediated apoptosis, natural killer cell-mediated apoptosis and in T-cell development.
别称	fas ligand, soluble Fas Ligand (sFasL), TNFSF6, CD95L, Apo I Ligand, APTL, APT1LG1, CD178, Fas-Lg, Tnfs, Tnlg1a, gld
蛋白编码	P41047
分子量	The protein has a calculated MW of 18.27 kDa. The protein migrates as 17-25 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>98% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce apoptosis in Jurkat cells. The ED $_{50}$ for this effect is <1 μg /mL.
内毒素检测	$<$ 0.1 EU per 1 μ g of the protein by the LAL method.
蛋白序列	QIANPSTPSEKKEPRSVAHLTGNPHSRSIPLEWEDTYGTALISGVKYKKGGLVINETGLYFVYSKV YFRGQSCNNQPLNHKVYMRNSKYPEDLVLMEEKRLNYCTTGQIWAHSSYLGAVFNLTSADH LYVNISQLSLINFEESKTFFGLYKL with polyhistidine tag at the N-terminus.
产品形式	The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than $10~\mu g/ml$.
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存 Initial Recoi

Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.


Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant mouse FasL

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491