# **CHAMOT**

重组鼠标肿瘤坏死因子超家族成员14 /Recombinant LIGHT, Mouse, Animal-Free


CM043-5MP

CM043-20MP

CM043-100MP

CM043-500MP

CM043-1000MP





# トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



### **Product Datasheet**

## 重组鼠标肿瘤坏死因子超家族成员14 /Recombinant LIGHT, Mouse, Animal-Free

| 产品编号 | CM043-5MP | CM043-20MP | CM043-100MP | CM043-500MP | CM043-<br>1000MP |
|------|-----------|------------|-------------|-------------|------------------|
| 规格   | 5 μg      | 20 μg      | 100 µg      | 500 μg      | 1 mg             |

### 产品简介

| 背景描述  | LIGHT, also known as TNFSF14 is a member of the TNF superfamily that produced by multiple immune cells such as activated T cells and immature dendritic cells (DCs). Mouse LIGHT shares 79% sequence homology with human LIGHT. LIGHT is a 29 kDa type II transmembrane protein, which serves as a ligand for both lymphotoxin $\beta$ receptor (LT $\beta$ R) and TNFSF signaling receptors (TNFRSF14/HVEM). Besides, LIGHT can amplify NF-kB signaling pathway in T cells when presencing anti-CD3 antibody treatment. Additionally, LIGHT is able to stimulate T cell proliferation and IFN expression via interacting with TNFRS13 /HVEM. |  |  |
|-------|---|--|--|
| 别称    | TNFSF14, HVEM-L, CD258, Ly113, LTg  |  |  |
| 蛋白编码  | Q9QYH9  |  |  |
| 分子量   | The protein has a calculated MW of 20.92 kDa. The protein migrates about 17 kDa under reducing condition (SDS-PAGE analysis).   |  |  |
| 表达系统  | Escherichia coli  |  |  |
| 纯度    | >98% as determined by SDS-PAGE.   |  |  |
| 生物活性  | Measure by its ability to induce cytotoxicity in HT-29 cells. The ED $_{50}$ for this effect is <2 $\mu g/mL$ .   |  |  |
| 内毒素检测 | $<$ 0.1 EU per 1 $\mu g$ of the protein by the LAL method.  |  |  |
| 蛋白序列  | MRLHQRLGDIVAHLPDGGKGSWEKLIQDQRSHQANPAAHLTGANASLIGIGGPLLWETRLGL AFLRGLTYHDGALVTMEPGYYYVYSKVQLSGVGCPQGLANGLPITHGLYKRTSRYPKELELLVS RRSPCGRANSSRVWWDSSFLGGVVHLEAGEEVVVRVPGNRLVRPRDGTRSYFGAFMV with polyhistidine tag at the C-terminus.   |  |  |
| 产品形式  | The protein was lyophilized from a 0.2 $\mu m$ filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.   |  |  |

### 产品使用

**1.** Before opening, centrifuge at 3000 rpm for 5 mins.

### 2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

### 3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than  $10~\mu g/ml$ .
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

### 产品储存/运输

储存

**Lyophilized Protein:** Store at -20°C for 1 year.


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

**运输** Blue Ice

### 实验数据展示



 ${\tt SDS-PAGE} \ analysis \ of \ recombinant \ mouse \ {\tt LIGHT}$ 

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491