# **CHAMOT**

### 重组人增殖诱导配体 /Recombinant APRIL,Human, Animal-Free

CM048-5HP

CM048-20HP

CM048-100HP

CM048-500HP

CM048-1000HP





# トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



### **Product Datasheet**

## 重组人增殖诱导配体 /Recombinant APRIL,Human,Animal-Free

产品编号	CM048-5HP	CM048-20HP	CM048-100HP	CM048-500HP	CM048-1000HP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

### 产品简介

背景描述	APRIL is a member of tumor necrosis factor ligand superfamily that expressed by stromal tissue, macrophages, and T cells. APRIL is a 27.4 kDa protein containing 250 residues, which plays a critical role in modulating tumor progression. Besides, APRIL has been demonstrated to involve in regulating B and T cell survival, proliferation and differentiation via binding with TNFRSF13B/TACI and TNFRSF17/BCMA.		
别称	A proliferation-inducing ligand, TNFSF13, CD256, TALL-2, TALL2, TNLG7B, TRDL-1, UNQ383/PRO715, ZTNF2		
蛋白编码	AAQ91388		
分子量	The protein has a calculated MW of 17.29 kDa. The protein migrates as 18 kDa under reducing condition (SDS-PAGE analysis).		
表达系统	Escherichia coli		
纯度	>98% as determined by SDS-PAGE.		
生物活性	Measured by its ability to induce cell death in Jurkat cells. The ED $_{50}$ for this effect is <6.0 $\mu g/mL$ .		
内毒素检测	$<$ 0.1 EU per 1 $\mu$ g of the protein by the LAL method.		
蛋白序列	MAVLTQKQKKQHSVLHLVPINATSKDDSDVTEVMWQPALRRGRGLQAQGYGVRIQDAGVYL LYSQVLFQDVTFTMGQVVSREGQGRQETLFRCIRSMPSHPDRAYNSCYSAGVFHLHQGDILS VIIPRARAKLNLSPHGTFLGFVKL with polyhistidine tag at the C-terminus.		
产品形式	The protein was lyophilized from a 0.2 $\mu m$ filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.		
产品应用	Cell Culture		

### 产品使用

**1.** Before opening, centrifuge at 3000 rpm for 5 mins.

### 2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

### 3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than  $10~\mu g/ml$ .
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

### 产品储存/运输

储存

**Lyophilized Protein:** Store at -20°C for 1 year.


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

**Extended Storage:** Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

**运输** Blue Ice

### 实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491