CHAMOT

重组人CD40配体 /Recombinant CD40L,Human,Animal-Free

CM053-5HP

CM053-20HP

CM053-100HP

CM053-500HP

CM053-1000HP





トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



Product Datasheet

重组人CD40配体 /Recombinant CD40L,Human,Animal-Free

产品编号	CM053-5HP	CM053-20HP	CM053-100HP	CM053-500HP	CM053-1000HP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

产品简介

背景描述	Human CD40L (CD154) is one member of TNF family which is expressed on T cells. It binds the CD40, which is on the antigen-presenting cells (ex: B cells, macrophages and monocytes) Human CD40L is a 21.6 kDa cytokine with 103 amino acid residues which is a transmembrane glycoprotein. Human CD40L plays an important role in the adaptive immune system which induces the B cell maturation and activation.
别称	CD40 ligand, soluble CD40 Ligand, TNFSF5, TRAP, CD154, Gp39, T-BAM, sCD40 Ligand
蛋白编码	P29965
分子量	The protein has a calculated MW of 16.99 kDa. The protein migrates as 16 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>95% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce IL-8 secretion in human PBMCs. The ED $_{50}$ for this effect is <5 ng/mL.
内毒素检测	$<$ 0.1 EU per 1 μ g of the protein by the LAL method.
蛋白序列	MQKGDQNPQIAAHVISEASSKTTSVLQWAEKGYYTMSNNLVTLENGKQLTVKRQGLYYIYAQ VTFCSNREASSQAPFIASLCLKSPGRFERILLRAANTHSSAKPCGQQSIHLGGVFELQPGASVFV NVTDPSQVSHGTGFTSFGLLKL with polyhistidine tag at the C-terminus
产品形式	The protein was lyophilized from a 0.2 μm filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than $10~\mu g/ml$.
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

储存

Lyophilized Protein: Store at -20°C for 1 year.


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输 Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491