

CHAMOT

**重组小鼠4-1BB配体 /Recombinant 4-1BBL,
Mouse, Animal-Free**

CM055-5MP

CM055-20MP

CM055-100MP

CM055-500MP

CM055-1000MP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组小鼠4-1BB配体 /Recombinant 4-1BBL, Mouse, Animal-Free

产品编号	CM055-5MP	CM055-20MP	CM055-100MP	CM055-500MP	CM055-1000MP
规格	5 µg	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述

4-1BB ligand (4-1BBL) is a type II transmembrane protein that is part of the tumor necrosis factor (TNF) ligand family. As an inducible co-stimulatory molecule, it presents on several antigen presenting cell (APC) types, including B cells, macrophages and DCs. The interactions between 4-1BB and 4-1BBL trigger pleiotropic effects on the immune response including antigen presenting process, proliferation of CD4 and CD8 positive T-cells, as well as cytokine secretion from T-cells through NFκB, c-Jun, and p38 downstream signal pathways activation. Therefore, 4-1BB and 4-1BBL are recently used for the immunotherapy of cancer.

别称

4-1BB ligand, 4-1BB, 4-1BB-L, 4-1BBL, AI848817, Cd137, Cd137l, Ly6, Ly63l, tumor necrosis factor (ligand) superfamily, member 9 , Tnfsf9

蛋白编码

P41274

分子量

The protein has a calculated MW of 23.9 kDa. The protein migrates as 24 kDa under reducing condition (SDS-PAGE analysis).

表达系统

Escherichia coli

纯度

>95% as determined by SDS-PAGE.

生物活性

Measure by its ability to induce IL-2 secretion in mouse T cells in the presence of the anti-CD3 antibody. The ED₅₀ for this effect is <0.05 µg/mL.

内毒素检测

<0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列

MRTEPRPALTITTSPNLGTRENNADQVTPVSHIGCPNTTQQGSPVFAKLLAKNQASLCNTTLN
WHSQDGAGSSYLSQGLRYEEDKKELVVDSPGLYYVFLELKLSPFTTNTGHKVQGWVSLVLQAK
PQVDDFDNLALTVELFPCSMENKLVDRWSQLLLLKAGHRLSVGLRAYLHGAQDAYRDWELS
YPNTTSFGLFLVKPDNPWE with polyhistidine tag at the C-terminus.

产品形式

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.

- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



SDS-PAGE analysis of recombinant mouse 41BBL

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491