# **CHAMOT**

# 重组人Fas配体 /Recombinant FasL, Human, Animal-Free

CM056-5HP

CM056-20HP

CM056-100HP

CM056-500HP

CM056-1000HP





# トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



# **Product Datasheet**

# 重组人Fas配体 /Recombinant FasL, Human, Animal-Free

| 产品编号 | CM056-5HP | CM056-20HP | CM056-100HP | CM056-500HP | CM056-1000HP |
|------|-----------|------------|-------------|-------------|--------------|
| 规格   | 5 μg      | 20 μg      | 100 μg      | 500 μg      | 1 mg         |

| 产品简介  |  |
|-------|--|
| 背景描述  | Fas Ligand (FasL) is a 17.34 kDa tumor necrosis factor with 152 amino acid residues. FasL is expressed from lymphoid tissue and secreted to blood. Binding to its receptor, TNFRSF6/FAS, leads to induce apoptotic signal into cells. Involved in cytotoxic T-cell-mediated apoptosis, natural killer cell-mediated apoptosis and in T-cell development. |
| 别称    | fas ligand, soluble Fas Ligand (sFasL), TNFSF6, CD95L, Apo I Ligand, APTL, APT1LG1, CD178  |
| 蛋白编码  | P48023   |
| 分子量   | The protein has a calculated MW of 29.51 kDa. The protein migrates as 30-35 kDa under reducing condition (SDS-PAGE analysis).  |
| 表达系统  | Escherichia coli   |
| 纯度    | >98% as determined by SDS-PAGE.  |
| 生物活性  | Measure by its ability to induce apoptosis in Jurkat cells. The ED $_{50}$ for this effect is <1 ng/mL. The specific activity of recombinant human FasL is > 1 x 10 $^{6}$ IU/mg.  |
| 内毒素检测 | $<$ 0.1 EU per 1 $\mu$ g of the protein by the LAL method.   |
| 蛋白序列  | QIGHPSPPPEKKELRKVAHLTGKSNSRSMPLEWEDTYGIVLLSGVKYKKGGLVINETGLYFVYSK VYFRGQSCNNLPLSHKVYMRNSKYPQDLVMMEGKMMSYCTTGQMWARSSYLGAVFNLTS ADHLYVNVSELSLVNFEESQTFFGLYKL with polyhistidine tag and sumo tag at the N-terminus   |
| 产品形式  | The protein was lyophilized from a 0.2 $\mu$ m filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.  |
| 产品应用  | Cell Culture   |

**1.** Before opening, centrifuge at 3000 rpm for 5 mins.

### 2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

### 3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than  $10~\mu g/ml$ .
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

# 产品储存/运输

**Lyophilized Protein:** Store at -20°C for 1 year.

储存 Initial Reconstitution


**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

**运输** Blue Ice

## 实验数据展示



SDS-PAGE analysis of recombinant human FasL

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491