# **CHAMOT**

重组人骨形态发生蛋白-4 /Recombinant BMP-4, Human, Animal-Free


CM065-5HP

CM065-20HP

CM065-100HP

CM065-500HP

CM065-1000HP





# トスツと

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示



### **Product Datasheet**

## 重组人骨形态发生蛋白-4 /Recombinant BMP-4, Human, Animal-Free

产品编号	CM065-5HP	CM065-20HP	CM065-100HP	CM065-500HP	CM065-1000HP
规格	5 μg	20 μg	100 μg	500 μg	1 mg

女	44	✓	
	ны	71	ı
	-		

背景描述	Bone Morphogenetic Protein-4 (BMP-4) is an extracellular multifunctional signaling cytokine that is also a member of the TGF- $\beta$ family. BMP-4 can bind with the TGF- $\beta$ receptor and trigger SMAD protein signal transduction. BMP-4 has a significant expression in the prostate, it plays a vital role in the development of mesoderm into bones and the development of the epidermis and Dorsal-Ventral axis.
别称	bone morphogenetic protein 4, BMP-2B, DVR4, BMP4
蛋白编码	P12644
分子量	The protein has a calculated MW of 12.88 kDa. The protein migrates as 12 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>95% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce alkaline phosphatase production by ATDC5 cells. The $ED_{50}$ for this effect is <0.58 ng/mL.
内毒素检测	$<$ 0.1 EU per 1 $\mu$ g of the protein by the LAL method.
蛋白序列	MKKNKNCRRHSLYVDFSDVGWNDWIVAPPGYQAFYCHGDCPFPLADHLNSTNHAIVQTLV NSVNSSIPKACCVPTELSAISMLYLDEYDKVVLKNYQEMVVEGCGCR with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 $\mu$ m filtered solution containing 1×PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

**1.** Before opening, centrifuge at 3000 rpm for 5 mins.

### 2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H<sub>2</sub>O to a concentration of 100-200 μg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

• Store at 2°C to 8°C for up to 1 week.

### 3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than  $10~\mu g/ml$ .
- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

### 产品储存/运输

**Lyophilized Protein:** Store at -20°C for 1 year.

储存 Initial Reconstitution:

**Initial Reconstitution:** Store at 2°C to 8°C for up to 1 week.


**Extended Storage:** Store at -20°C or -80°C for 3 to 6 months with a carrier protein or

stabilizer.

运输

Blue Ice

### 实验数据展示



SDS-PAGE analysis of recombinant human BMP-4

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491