

CHAMOT

**重组人干扰素omega /Recombinant IFN omega,
Human, Animal-Free**

CM082-20HP

CM082-100HP

CM082-500HP

CM082-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组人干扰素omega /Recombinant IFN omega, Human, Animal-Free

产品编号	CM082-20HP	CM082-100HP	CM082-500HP	CM082-1000HP
规格	20 µg	100 µg	500 µg	1 mg

产品简介

背景描述	Interferon Omega (IFN- ω) is a 20.12 kDa member of type I IFN family with 173 amino acid residues. IL-28B is expressed by epithelial tissues. IFN- ω with antiviral, antitumor activity and regulating the innate immune response. Able to activate P13K/Akt signaling pathway via binding its receptor IFNAR in cells.
别称	interferon omega, IFN alpha II-1, IFNW1
蛋白编码	P05000
分子量	The protein has a calculated MW of 20.93 kDa. The protein migrates as 20 kDa under reducing condition (SDS-PAGE analysis).
表达系统	Escherichia coli
纯度	>95% as determined by SDS-PAGE.
生物活性	Measure by its ability to induce cytotoxicity in TF-1 cells. The ED ₅₀ for this effect is <0.02 ng/mL. The specific activity of recombinant human IFN omega is approximately >5 x10 ⁷ IU/ mg.
内毒素检测	<0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列	MCDLPQNHGLLSRNTLVLLHQMRRISPFLCLKDRRDFRFPQEMVKGSQQLQKAHVMSVLHEMLQQIFSLFHTERSSAAWNMTLLDQLHTGLHQQQLQHLETCLLQVVGEGESAGAISSPALTLLRRYFQGIRVYLKEKKYSDCAWEVVRMEIMKSLFLSTNMQERLRSKDRDLGSS with polyhistidine tag at the C-terminus.
产品形式	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.
产品应用	Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.

- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491