

CHAMOT

重组小鼠趋化因子CXCL7 /Recombinant CXCL7(48-109), Mouse, Animal-Free

CM088-5MP

CM088-20MP

CM088-100MP

CM088-500MP

CM088-1000MP



CHAMOT

乔默®生物

Specialize In Cytokines



CONTENT

1 产品简介

2 产品使用

3 产品储存/运输

4 实验数据展示

重组小鼠趋化因子CXCL7 /Recombinant CXCL7(48-109), Mouse, Animal-Free

| 产品编号 | CM088-5MP | CM088-20MP | CM088-100MP | CM088-500MP | CM088-1000MP |
|------|-----------|------------|-------------|-------------|--------------|
| 规格 | 5 µg | 20 µg | 100 µg | 500 µg | 1 mg |

产品简介

背景描述

C-X-C motif chemokine 7 (CXCL7) also named Pro-Platelet basic protein (PPBP), which is a chemokine of the intercrine alpha family. CXCL7 is a 8.2 kDa protein containing 74 amino acid residues. CXCL7 is expressed by the platelets, which are activated. During vascular injury, CXCL7 controls the glucose metabolism, mitogenesis and neutrophil recruitment by the interaction with CXCR2.

别称

C-X-C motif chemokine 7, NAP-2, PPBP, CT, CTA, CTAP3, CTAPIII, LA-PF4, LDG, LDGF, MDGF, NAP-2-L1, Scyb, Scyb7, TGB, TGB1, THBGB1, b-TG1, beta-T, beta-TG

蛋白编码

NP_076274

分子量

The protein has a calculated MW of 7.57 kDa. The protein migrates below 11 kDa under reducing condition (SDS-PAGE analysis).

表达系统

Escherichia coli

纯度

>98% as determined by SDS-PAGE.

生物活性

Measure by its ability to chemoattract human THP1 cells using a concentration range of 50-100 ng/mL.

内毒素检测

<0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列

IELRCRCTNTISGIPFNSISLVNVYRPGVHCADVEVIATLKNGQKTCLDPNAPGVKRIVMKI with polyhistidine tag at the N-terminus.

产品形式

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

产品应用

Cell Culture

产品使用

1. Before opening, centrifuge at 3000 rpm for 5 mins.

2.Initial Reconstitution

- Reconstitute the Lyophilized Protein in sterile H₂O to a concentration of 100-200 µg/mL.
- Then, incubate it at room temperature for at least 20 mins to ensure sufficient dissolution.

Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

- Store at 2°C to 8°C for up to 1 week.

3.Extended Storage

- After the initial reconstitution, further dilute the reconstituted protein in a buffer containing a carrier protein or stabilizer (e.g., 0.1% BSA, 10% FBS, 5% HSA, or 5% trehalose solution). The final concentration is not less than 10 µg/ml.

- Prepare aliquots (≥20 µl).
- Store at -20°C or -80°C for 3 to 6 months.

Avoid repeated freeze-thaw cycles.

产品储存/运输

Lyophilized Protein: Store at -20°C for 1 year.

储存


Initial Reconstitution: Store at 2°C to 8°C for up to 1 week.

Extended Storage: Store at -20°C or -80°C for 3 to 6 months with a carrier protein or stabilizer.

运输

Blue Ice

实验数据展示



For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491